

NYRx Drug Class Coverage Overview: HIV Post-Exposure Prophylaxis Agents

NYRx Drugs Covered for Post-Exposure Prophylaxis to Prevent HIV Infection

As outlined in the New York State Department of Health AIDS Institute [Post-Exposure Prophylaxis \(PEP\) to Prevent HIV Infection Clinical Guidelines](#), the following drugs can be used for HIV PEP:

- Cimduo (lamivudine / tenofovir disoproxil fumarate)
- Emtriva (emtricitabine)
- Epivir, Epivir HBV (lamivudine)
- Isentress (raltegravir)
- Kaletra (lopinavir / ritonavir)
- Lexiva (fosamprenavir)
- Norvir (ritonavir)
- Retrovir (zidovudine)
- Reyataz (atazanavir)
- Stribild (elvitegravir, cobicistat, emtricitabine, and tenofovir disoproxil fumarate)
- Tivicay (dolutegravir)
- Truvada (emtricitabine / tenofovir)
- Viread (tenofovir)

What Pharmacy Providers Need to Do

Pharmacy providers should bill PEP agents for a 28-day supply.

What Prescribers Need to Do

Prescribers should write a prescription for a 28-day supply of PEP. A prior authorization is not required if the prescription is written for a 28-day supply, there is no diagnosis of HIV and no HIV drugs in the members history in the last 120 days.

Resources

- [NYRx Education & Outreach Website](#)
- [NYRx Prior Authorization Submission Guide](#)
- [New York State Medicaid Drug Utilization Program](#)

Contact Information

The NYRx Education & Outreach Call Center is available by phone at 1-833-967-7310 or by email at NYRxEO@primetherapeutics.com from 8:00 AM to 5:00 PM ET, Monday through Friday, excluding holidays.

The NYRx Education & Outreach team hosts virtual office hours every week for stakeholders to ask questions related to NYRx and care coordination. Visit the [NYRx Education & Outreach website](#) for more information.