

# NYRx, the New York Medicaid Pharmacy Program

## OVERVIEW OF CONTENTS

### Preferred Drug Program (PDP) (Pages 4–61)

The PDP promotes the use of less expensive, equally effective drugs when medically appropriate through a Preferred Drug List (PDL). All drugs currently covered by NYRx, the Medicaid Pharmacy Program, remain available under the PDP and the determination of preferred and non-preferred drugs does not prohibit a prescriber from obtaining any of the medications covered under Medicaid.

- Non-preferred drugs in these classes require prior authorization (PA), unless indicated otherwise.
- Preferred drugs that require prior authorization are indicated by footnote.
- Specific Clinical, Frequency/Quantity/Duration, Step Therapy criteria is listed in column at the right.

### Clinical Drug Review Program (CDRP) (Page 62)

The CDRP is aimed at ensuring specific drugs are utilized in a medically appropriate manner. Under the CDRP, certain drugs require prior authorization because there may be specific safety issues, public health concerns, the potential for fraud and abuse, or the potential for significant overuse and misuse.

### Drug Utilization Review (DUR) Program (Pages 63–77)

The DUR helps to ensure that prescriptions for outpatient drugs are appropriate, medically necessary, and not likely to result in adverse medical consequences. This program uses professional medical protocols and computer technology and claims processing to assist in the management of data regarding the prescribing and dispensing of prescriptions. Frequency/Quantity/Duration (F/Q/D) Program and Step Therapy parameters are implemented to ensure clinically appropriate and cost-effective use of these drugs and drug classes.

### Medication Assisted Treatment (MAT) Formulary (Page 78)

Prior authorization will not be required for medications used for the treatment of substance use disorder prescribed according to generally accepted national professional guidelines for the treatment of a substance use disorder.

### Brand Less Than Generic (BLTG) Program (Pages 79–80)

The Brand Less Than Generic Program is a cost containment initiative which promotes the use of certain multi-source brand name drugs when the cost of the brand name drug is less expensive than the generic equivalent. This program is in conformance with State Education Law, which intends that patients receive the lower cost alternative.

### Mandatory Generic Drug Program (Page 81)

State law excludes Medicaid coverage of brand name drugs that have a Federal Food and Drug Administration (FDA) approved A-rated generic equivalent unless a prior authorization is obtained. Drugs subject to the Preferred Drug Program (PDP), Clinical Drug Review Program (CDRP), and/or the Brand Less Than Generic (BLTG) Program are not subject to the Mandatory Generic Program.

For more information on NYRx, the Medicaid Pharmacy Program: [http://www.health.ny.gov/health\\_care/medicaid/program/pharmacy.htm](http://www.health.ny.gov/health_care/medicaid/program/pharmacy.htm)

To contact the NYRx Clinical Call Center please call 1-877-309-9493

To download a copy of the Prior Authorization fax form go to [https://newyork.fhsc.com/providers/PA\\_forms.asp](https://newyork.fhsc.com/providers/PA_forms.asp)

Disclaimer: Branded generics are included with the single generic name listing; they are not listed as separate agents.

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

### Dose Optimization Program (Pages 82–87)

Dose optimization can reduce prescription costs by reducing the number of pills a patient needs to take each day. The Department has identified drugs to be included in this program, the majority of which have FDA approval for once-a-day dosing, have multiple strengths available in correlating increments at similar costs and are currently being utilized above the recommended dosing frequency.

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

## PREFERRED DRUG LIST – TABLE OF CONTENTS

I. ANALGESICS .....	4
II. ANTI-INFECTIVES.....	8
III. CARDIOVASCULAR .....	11
IV. CENTRAL NERVOUS SYSTEM .....	17
V. DERMATOLOGIC AGENTS.....	28
VI. ENDOCRINE AND METABOLIC AGENTS.....	34
VII. GASTROINTESTINAL.....	40
VIII. HEMATOLOGICAL AGENTS .....	43
IX. IMMUNOLOGIC AGENTS .....	46
X. MISCELLANEOUS AGENTS .....	48
XI. MUSCULOSKELETAL AGENTS.....	49
XII. OPHTHALMICS.....	50
XIII. OTICS.....	54
XIV. RENAL AND GENITOURINARY .....	55
XV. RESPIRATORY .....	57

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>I. Analgesics</b>		
<b>Non-Steroidal Anti-Inflammatory Drugs (NSAIDS)</b>		
diclofenac 1% topical gel diclofenac sodium oral ibuprofen Rx tablet, suspension ibuprofen OTC suspension indomethacin ketorolac meloxicam tablet naproxen tablet piroxicam sulindac	Arthrotec® Celebrex® <small>CC</small> celecoxib <small>CC</small> Daypro® diclofenac epolamine patch (gen Flector®) diclofenac capsule diclofenac/misoprostol diclofenac potassium diclofenac potassium (gen Cambia®) diclofenac sodium ER diclofenac topical soln (gen Pennsaid®) diflunisal Duexis® Elyxyb™ <small>F/Q/D</small> etodolac etodolac ER Feldene® fenoprofen Flector® patch flurbiprofen ibuprofen/famotidine (gen Duexis®) indomethacin ER ketoprofen ketoprofen ER ketorolac nasal spray (gen Sprix®) Licart™ meclofenamate mefenamic acid meloxicam capsule (gen Vivlodex®) Mobic®	<b>CLINICAL CRITERIA (CC)</b> <ul style="list-style-type: none"> <li>• <b>Celebrex® (celecoxib)</b> – one of the following criteria will not require PA <ul style="list-style-type: none"> <li>– Over the age of 65 years</li> <li>– Concurrent use of an anticoagulant agent</li> <li>– History of GI Bleed/Ulcer or Peptic Ulcer Disease</li> </ul> </li> </ul> <b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b> <ul style="list-style-type: none"> <li>• Elyxyb™ (celecoxib) – 4.8 mL bottle (120 mg) maximum quantity: 9 / 30 days</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>I. Analgesics</b>		
	nabumetone Nalfon® Naprelan® naproxen susp naproxen CR naproxen EC naproxen-esomeprazole naproxen sodium oxaprozin Pennsaid® Relafen® DS tolmetin Vimovo®	
<b>Opioids – Long-Acting <sup>CC</sup></b>		
buprenorphine patch  fentanyl patch (12 mcg, 25 mcg, 50 mcg, 75 mcg, 100 mcg)  morphine sulfate ER tablet	Belbuca® Butrans® ConZip® <sup>ST</sup>  fentanyl patch (37.5 mcg, 62.5 mcg, 87.5 mcg)  hydrocodone ER  hydrocodone ER (gen Hysingla ER)  hydromorphone ER  Hysingla® ER  morphine ER capsule (gen Avinza)  morphine ER capsule (gen Kadian)  MS Contin®  Nucynta® ER <sup>ST</sup>  oxycodone ER  Oxycontin®  oxymorphone ER  tramadol ER <sup>ST</sup>  Xtampza® ER	<b>CLINICAL CRITERIA (CC) *</b> <ul style="list-style-type: none"> <li>• Limited to a total of 4 opioid prescriptions every 30 days; Exemption for diagnosis of cancer or sickle cell disease</li> <li>• PA required for initiation of opioid therapy for patients on established opioid dependence therapy</li> <li>• PA required for use if <math>\geq</math> 90 MME (MME = morphine milligram equivalents) of opioid per day for management of non-acute pain (pain lasting <math>&gt;</math> 7 days)</li> <li>• PA required for initiation of long-acting opioid therapy in opioid-naïve patients.</li> <li>• PA required for any additional long-acting opioid prescription for patients currently on long-acting opioid therapy.</li> <li>• PA required for initiation of opioid therapy in patients currently on benzodiazepine therapy</li> <li>• PA required for any codeine- or tramadol-containing products in pts <math>&lt;</math> 12 years</li> <li>• PA required for initiation of opioid therapy for patients on established CNS stimulant therapy</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>I. Analgesics</b>		
		<p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>• <b>Nucynta® ER (tapentadol ER)</b>: Trial with tapentadol IR before tapentadol ER for patients who are naïve to a long-acting opioid</li> <li>• <b>Tramadol ER (tramadol naïve patients)</b>: Attempt treatment with IR formulations before the following ER formulations: ConZip®, tramadol ER</li> </ul> <p>*Exemption from requirements for diagnosis of cancer, sickle cell disease, or hospice care.</p>
<b>Opioids – Short-Acting <sup>CC</sup></b>		
butalbital/APAP/caffeine/codeine codeine codeine/APAP hydrocodone/APAP hydrocodone/ibuprofen morphine IR oxycodone/APAP tramadol tablet	Apadaz® benzhydrocodone/APAP butalbital compound/codeine butorphanol nasal spray dihydrocodeine/APAP/caffeine Dilauidid® hydromorphone levorphanol meperidine Nalocet® Nucynta® <sup>ST</sup> oxycodone oxymorphone pentazocine/naloxone Percocet® Prolate® solution, tablet Roxicodone® Seglentis® tramadol solution tramadol/APAP	<p><b>CLINICAL CRITERIA (CC) *</b></p> <ul style="list-style-type: none"> <li>• Limited to a total of 4 opioid prescriptions every 30 days.</li> <li>• Initial prescription for opioid-naïve patients limited to a 7-day supply.</li> <li>• PA required for initiation of opioid therapy for patients on established opioid dependence therapy.</li> <li>• PA required for use if ≥ 90 MME of opioid per day for management of non-acute pain (&gt; 7 days) <ul style="list-style-type: none"> <li>– Exception for diagnosis of cancer or sickle cell disease, or hospice program</li> </ul> </li> <li>• PA is required for opioid-naïve patients for prescription requests ≥ 50 MME per day.</li> <li>• PA required for continuation of opioid therapy beyond an initial 7-day supply in patients established on gabapentin or pregabalin</li> <li>• PA required for initiation of opioid therapy in patients currently on benzodiazepine therapy</li> <li>• PA required for any codeine- or tramadol-containing products in pts &lt; 12 years</li> <li>• PA required for initiation of opioid therapy for patients on &gt; 7 days established CNS stimulant therapy</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>• <b>Nucynta® (tapentadol IR)</b> – Trial with tramadol and 1 preferred opioid before tapentadol immediate-release (IR)</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>I. Analgesics</b>		
		<ul style="list-style-type: none"> <li>For Non-opioid Pain management alternatives please visit:  <a href="https://health.ny.gov/health_care/medicaid/program/opioid_management/docs/non_opioid_alternatives_to_pain_management.pdf">https://health.ny.gov/health_care/medicaid/program/opioid_management/docs/non_opioid_alternatives_to_pain_management.pdf</a> </li> </ul> <p>*Exemptions from requirements for diagnosis of cancer, sickle cell disease, or hospice care</p>

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>II. Anti-Infectives</b>		
<b>Antibiotics – Inhaled CC, F/Q/D</b>		
Bethkis® <small>BLTG</small> Cayston® Kitabis® Pak <small>BLTG</small> TOBI Podhaler™ tobramycin (gen TOBI®) solution	TOBI® solution tobramycin (gen Bethkis®, Kitabis®) solution	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>Aztreonam (Cayston) <ul style="list-style-type: none"> <li>3 ampules (3 mL) per day</li> <li>84 ampules (84 mL) per 56-day regimen (28 days on, 28 days off)</li> </ul> </li> <li>Tobramycin inhalation solution (Bethkis, TOBI, Kitabis Pak) <ul style="list-style-type: none"> <li>2 ampules (8 mL Bethkis, 10 mL TOBI, Kitabis Pak) per day</li> <li>56 ampules (224 mL Bethkis, 280 mL TOBI, Kitabis Pak) per 56 day regimen (28 days on-28 days off)</li> </ul> </li> <li>Tobramycin capsules with inhalation powder (TOBI Podhaler) <ul style="list-style-type: none"> <li>8 capsules per day 224 capsules per 56-day regimen (28 days on-28 days off)</li> </ul> </li> </ul>
<b>Anti-Fungals – Oral for Onychomycosis</b>		
griseofulvin suspension, ultramicrogranulated terbinafine tablet	griseofulvin tablet itraconazole itraconazole solution (gen Sporanox) Sporanox®	
<b>Anti-Virals – Oral</b>		
acyclovir valacyclovir	famciclovir Valtrex® Zovirax®	
<b>Cephalosporins – Third Generation</b>		
cefdinir	cefixime cefpodoxime Suprax®	
<b>Fluoroquinolones – Oral</b>		
ciprofloxacin suspension, tablet	Baxdela®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>II. Anti-Infectives</b>		
levofloxacin tablet	Cipro® suspension, tablet levofloxacin solution moxifloxacin ofloxacin tablet	
<b>Hepatitis B Agents</b>		
adefovir dipivoxil Baraclude® solution entecavir Epivir-HBV® solution lamivudine HBV	Baraclude® tablet Epivir-HBV® tablet Hepsera® Vemlidy®	
<b>Hepatitis C Agents – Direct Acting Antivirals</b>		
Mavyret™ CC, F/Q/D  ribavirin  sofosbuvir/velpatasvir (gen Epclusa®) CC, F/Q/D  Vosevi® CC, F/Q/D	Epclusa® CC, F/Q/D  Harvoni® CC, F/Q/D  ledipasvir/sofosbuvir (gen Harvoni®) CC, F/Q/D  Sovaldi® CC, F/Q/D  Viekira Pak® CC, F/Q/D  Zepatier® CC, F/Q/D	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> <li>For patients being retreated require confirmation of patient readiness and adherence <ul style="list-style-type: none"> <li>Evaluation by using scales or assessment tools readily to determine a patient's readiness to initiate HCV treatment, specifically drug and alcohol abuse potential. Assessment tools are available to healthcare practitioners at: <a href="https://www.drugabuse.gov/nidamed-medical-health-professionals/screening-tools-resources/chart-screening-tools">https://www.drugabuse.gov/nidamed-medical-health-professionals/screening-tools-resources/chart-screening-tools</a></li> <li>OR <a href="https://prepc.org/">https://prepc.org/</a>.</li> </ul> </li> <li>The optional Hepatitis C Worksheet can be accessed at: <a href="https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Worksheet_Prescribers_HepC.pdf">https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Worksheet_Prescribers_HepC.pdf</a></li> </ul>
<b>Tetracyclines</b>		
demeclocycline  doxycycline hyolate  minocycline capsule  tetracycline	Doryx® ST, F/Q/D  Doryx MPC® ST, F/Q/D  doxycycline hyolate DR ST, F/Q/D  doxycycline monohydrate  minocycline tablet  minocycline ER tablet	<p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>Trial of doxycycline IR before progressing to doxycycline DR</li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>doxycycline DR (Doryx®): <ul style="list-style-type: none"> <li>Maximum 28 tablets/capsules per fill</li> </ul> </li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>II. Anti-Infectives</b>		
	Minolira ER™ Nuzyra™ Solodyn® Vibramycin®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>III. Cardiovascular</b>		
<b>Angiotensin Converting Enzyme Inhibitors (ACEIs)</b>		
benazepril enalapril lisinopril ramipril	Accupril® Altace® captopril enalapril (gen Epaned®) Epaned® fosinopril Lotensin® moexipril perindopril Qbrelis™ quinapril trandolapril Vasotec® Zestril®	
<b>ACE Inhibitor Combinations</b>		
benazepril/amlodipine benazepril/HCTZ captopril/HCTZ enalapril/HCTZ lisinopril/HCTZ Lotrel® trandolapril/verapamil ER	Accuretic® fosinopril/HCTZ Lotensin HCT® quinapril/HCTZ Vaseretic® Zestoretic®	

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>III. Cardiovascular</b>		
<b>Angiotensin Receptor Blockers (ARBs)</b>		
irbesartan <sup>1</sup> losartan olmesartan <sup>1</sup> telmisartan <sup>1</sup> valsartan tablet	Atacand® Avapro® Benicar® <u>DO</u> candesartan Cozaar® Diovan® <u>DO</u> , <sup>2</sup> Edarbi® eprosartan Micardis® <u>DO</u>	<b>DOSE OPTIMIZATION (DO)</b> <ul style="list-style-type: none"><li>See Dose Optimization Chart for affected drugs and strengths</li></ul>
<b>Antianginals and Anti-Ischemics</b>		
ranolazine	Aspruzyo Sprinkle™ Ranexa®	
<b>ARBs Combinations</b>		
Entresto® Exforge HCT® irbesartan/HCTZ <sup>1</sup> losartan/HCTZ olmesartan/amlodipine <sup>1</sup> olmesartan/HCTZ <sup>1</sup> telmisartan/HCTZ <sup>1</sup> valsartan/amlodipine valsartan/amlodipine/HCTZ valsartan/HCTZ	Atacand HCT® Avalide® Azor® Benicar HCT® <u>DO</u> candesartan/HCTZ Diovan HCT® <u>DO</u> Edarbyclor® <u>DO</u> Exforge® <u>DO</u> Hyzaar® Micardis HCT® <u>DO</u> olmesartan/amlodipine/HCTZ telmisartan/amlodipine Tribenzor®	<b>DOSE OPTIMIZATION (DO)</b> <ul style="list-style-type: none"><li>See Dose Optimization Chart for affected drugs and strengths</li></ul>

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf)

12

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>III. Cardiovascular</b>		
<b>Beta Blockers</b>		
atenolol carvedilol labetalol metoprolol succ. XL metoprolol tartrate propranolol tablet	acebutolol betaxolol bisoprolol Bystolic® <small>DO</small> carvedilol ER Coreg® Coreg CR® <small>DO</small> Corgard® Inderal LA® Inderal XL® InnoPran XL® Kapspargo™ Sprinkle Lopressor® nadolol <small>DO</small> nebivolol (gen Bystolic®) pindolol propranolol solution propranolol ER/SA Tenormin® timolol Toprol XL® <small>DO</small>	<b>DOSE OPTIMIZATION (DO)</b> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected drugs and strengths</li> </ul>
<b>Beta Blockers / Diuretics</b>		
atenolol/chlorthalidone bisoprolol/HCTZ propranolol/HCTZ	metoprolol tartrate/ HCTZ Tenoretic® Ziac®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>III. Cardiovascular</b>		
<b>Calcium Channel Blockers (Dihydropyridine)</b>		
amlodipine felodipine ER isradipine nicardipine HCl nifedipine nifedipine ER/SA	Katerzia™ levamldipine nisoldipine Norliqva® Norvasc® Procardia XL® Sular®	
<b>Cholesterol Absorption Inhibitors</b>		
cholestyramine cholestyramine light Colestid® tablet colestipol tablet ezetimibe	colesevelam Colestid granules, packet colestipol granules, packet Questran® Questran Light® Welchol® Zetia®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>III. Cardiovascular</b>		
<b>HMG-CoA Reductase Inhibitors (Statins)</b>		
atorvastatin lovastatin pravastatin rosuvastatin simvastatin		
	Altoprev® Atorvaliq® atorvastatin/amlodipine Caduet® Crestor® <small>DO</small> Ezallor™ Sprinkle ezetimibe/simvastatin fluvastatin fluvastatin ER Lescol XL® Lipitor® Livalo® Vytorin® Zocor® Zypitamag™	<b>DOSE OPTIMIZATION (DO)</b> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected drugs and strengths</li> </ul>
<b>Phosphodiesterase Type-5 (PDE-5) Inhibitors for PAH <small>CC</small></b>		
sildenafil tadalafil	Adcirca® Liqrev® Revatio® Tadliq®	<b>CLINICAL CRITERIA</b> <ul style="list-style-type: none"> <li>All prescriptions for <b>Adcirca®</b>, <b>tadalafil</b>, <b>Revatio®</b>, and <b>sildenafil</b> must have PA</li> <li>Prescribers or their authorized agents are required to respond to a series of questions that identify prescriber, patient, and reason for prescribing drug</li> <li>Please be prepared to fax clinical documentation upon request</li> <li>Prescriptions can be written for a 30-day supply with up to 11 refills</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>III. Cardiovascular</b>		
<b>Pulmonary Arterial Hypertension (PAH) Agents, Other – Oral</b>		
ambrisentan (gen Letairis) bosentan tablets (gen Tracleer®)	Adempas® Letairis® Opsumit® Orenitram® ER tablet, dosepack Tracleer® tablet for suspension, tablet Tyvaso DPI™ Uptravi®	
<b>Triglyceride Lowering Agents</b>		
fenofibrate tablet (gen Tricor®) fenofibrate capsule (gen Lofibra®) fenofibric acid capsule (gen Trilipix®) gemfibrozil omega-3 ethyl ester (gen Lovaza®) <small>F/Q/D</small> Vascepa® <small>F/Q/D, BLTG, 1</small>	Antara® fenofibrate caps (gen Lipofen®) fenofibrate micronized capsule (gen Antara®) fenofibrate tablet (gen Fenoglide®) fenofibric acid tablet (gen Fibrincor®) Fenoglide® icosapent (gen Vascepa®) <small>F/Q/D</small> Lipofen® Lopid® Lovaza® <small>F/Q/D</small> Tricor® Trilipix®	<b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b> <ul style="list-style-type: none"> <li>Lovaza® (omega-3-acid ethyl-esters) and Vascepa® (icosapent ethyl) – Required dosage equal to 4 grams per day</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
<b>Alzheimer's Agents</b>		
donepezil 5 mg, 10 mg galantamine galantamine ER memantine Namenda® rivastigmine	Adlyary® Aricept® donepezil 23 mg Exelon® memantine ER <small>CC, ST</small> Namenda XR® <small>CC, ST</small> Namzaric® <small>CC, ST</small> Razadyne ER®	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Memantine extended-release containing products (Namenda XR® and Namzaric®) – Require confirmation of diagnosis of dementia or Alzheimer's disease</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>Memantine extended-release containing products (Namenda XR® and Namzaric®) – Require trial with memantine immediate-release Namenda®</li> </ul>
<b>Anticonvulsants – Carbamazepine Derivatives</b>		
carbamazepine chewable, tablet carbamazepine ER capsule Equetro® oxcarbazepine tablet Tegretol® suspension <small>BLTG</small> Tegretol XR® <small>BLTG</small> Trileptal® suspension <small>BLTG</small>	Aptiom® <small>CC, DO</small> carbamazepine suspension <small>CC</small> carbamazepine XR tablet Carbatrol® <small>CC</small> oxcarbazepine suspension Oxtellar XR® <small>CC, DO</small> Tegretol® tablet <small>CC</small> Trileptal® tablet <small>CC</small>	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Clinical editing will allow patients currently stabilized on a non-preferred agent to continue to receive that agent without PA</li> </ul> <p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected drugs and strengths</li> </ul>
<b>Anticonvulsants – Other</b>		
clobazam tablet <small>ST, CC</small> gabapentin capsule, solution, tablet <small>F/Q/D, CC</small> lacosamide tablet, solution <sup>1</sup> lamotrigine tablet, chew levetiracetam levetiracetam ER Lyrica® capsule <small>DO, F/Q/D, CC</small> pregabalin capsule <small>DO, F/Q/D, CC</small> tiagabine topiramate <small>CC</small>	Banzel® Briviact® clobazam suspension <small>ST</small> Diacomit® <small>CC</small> Elepsia® XR Epidiolex® <small>CC</small> Eprontia™ <small>CC</small> felbamate Felbatol® Fintepla® Fycompa® <small>DO</small>	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected drugs and strengths</li> </ul> <p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Clinical editing will allow patients currently stabilized on a non-preferred agent to continue to receive that agent without PA</li> <li><b>Cannabidiol extract (Epidiolex®)</b> – Confirm diagnosis of FDA-approved or compendia-supported indication, or; Institutional Review Board (IRB) approval with signed consent form</li> <li><b>Lyrica®/Lyrica® CR (pregabalin)</b> – PA required for the initiation of pregabalin at &gt; 150 mg per day in patients currently on an opioid at &gt; 50 MME per day</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
zonisamide	Gabitril® Keppra® Keppra XR® Lamictal® tablet, chew, dosepak Lamictal® ODT tablet, dosepak Lamictal® XR <small>DO</small> tablet, dosepak lamotrigine dosepak lamotrigine ER lamotrigine ODT dosepak Lyrica® solution <small>DO, F/Q/D</small> Lyrica® CR <small>F/Q/D, CC</small> Neurontin® <small>F/Q/D, CC</small> Onfi® <small>ST, CC</small> pregabalin solution <small>DO, F/Q/D, CC</small> pregabalin ER (gen Lyrica® CR) <small>F/Q/D, CC</small> Qudexy® XR <small>CC</small> rufinamide (gen Banzel®) Sabril® Spritam® Sympazan® film <small>ST, CC</small> Topamax® <small>CC</small> topiramate ER <small>CC, DO</small> (gen Qudexy® XR) topiramate ER <small>CC</small> (gen Trokendi XR®) Trokendi XR® <small>CC, DO</small> vigabatrin Vimpat® Xcopri® Zonisade™ Ztalmy®	<ul style="list-style-type: none"> <li>• <b>Neurontin® (gabapentin)</b> – PA required for initiation of gabapentin at &gt; 900 mg per day in patients currently on an opioid at &gt; 50 MME per day</li> <li>• <b>Stiripentol (Diacomit®)</b> – Require diagnosis of FDA-approved or compendia-supported indication, or; Institutional Review Board (IRB) approval with signed consent form</li> <li>• <b>Topiramate IR/ER (Eprontia™, Qudexy® XR, Topamax®, Trokendi XR™)</b> – Require confirmation of FDA-approved, compendia-supported, or Medicaid covered diagnosis</li> <li>• <b>Onfi®/Sympazan® (clobazam):</b> <ul style="list-style-type: none"> <li>– Require confirmation of FDA-approved or compendia-supported use</li> <li>– PA required for initiation of clobazam therapy in patients currently on opioid or oral buprenorphine therapy</li> <li>– PA required for any clobazam prescription in patients currently on benzodiazepine therapy</li> </ul> </li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>• <b>Eprontia™ (topiramate)</b> – Maximum quantity: 473 mL per month</li> <li>• <b>Lyrica®/Lyrica® CR (pregabalin)</b> – Maximum daily dose of IR: 600 mg per day, and ER: 660 mg per day</li> <li>• <b>Neurontin® (gabapentin)</b> – Maximum daily dose of 3,600 mg per day</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>• <b>Onfi®/Sympazan® (clobazam)</b> – Requires a trial with an SSRI or SNRI for treatment of anxiety</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters																				
<b>IV. Central Nervous System</b>																						
<b>Antimigraine Agents, Other <small>ST, F/Q/D</small></b>																						
Ajovy® Emgality® Nurtec™ ODT	Aimovig® Emgality® 100mg syringe Qulipta™ Revvow™ Ubrelvy™ Zavzpret™	<p><b>STEP THERAPY (ST)</b></p> <p><b>Acute treatment of migraine</b></p> <ul style="list-style-type: none"> <li>• Trial of a product from the Antimigraine Agents-Triptan class</li> </ul> <p><b>Prevention of migraine</b></p> <ul style="list-style-type: none"> <li>• Trial of 2 FDA approved or compendia supported migraine prevention products from other drug classes</li> </ul> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="background-color: #0070C0; color: white;">Agent</th><th style="background-color: #0070C0; color: white;">F/Q/D</th></tr> </thead> <tbody> <tr> <td>Aimovig</td><td>1 syringe/30 days</td></tr> <tr> <td>Emgality 120 mg</td><td>2 syringes/30 days</td></tr> <tr> <td>Emgality 100 mg</td><td>3 syringes/30 days</td></tr> <tr> <td>Ajovy</td><td>3 syringes/90 days</td></tr> <tr> <td>Revvow</td><td>8 units/30 days</td></tr> <tr> <td>Ubrelvy</td><td>16 units/30 days</td></tr> <tr> <td>Nurtec™ ODT</td><td>24 units/40 days</td></tr> <tr> <td>Qulipta</td><td>30 units/30 days</td></tr> <tr> <td>Zavzpret®</td><td>8 units/30 days</td></tr> </tbody> </table>	Agent	F/Q/D	Aimovig	1 syringe/30 days	Emgality 120 mg	2 syringes/30 days	Emgality 100 mg	3 syringes/30 days	Ajovy	3 syringes/90 days	Revvow	8 units/30 days	Ubrelvy	16 units/30 days	Nurtec™ ODT	24 units/40 days	Qulipta	30 units/30 days	Zavzpret®	8 units/30 days
Agent	F/Q/D																					
Aimovig	1 syringe/30 days																					
Emgality 120 mg	2 syringes/30 days																					
Emgality 100 mg	3 syringes/30 days																					
Ajovy	3 syringes/90 days																					
Revvow	8 units/30 days																					
Ubrelvy	16 units/30 days																					
Nurtec™ ODT	24 units/40 days																					
Qulipta	30 units/30 days																					
Zavzpret®	8 units/30 days																					
<b>Antimigraine Agents – Triptans</b>																						
rizatriptan <small>F/Q/D</small> sumatriptan <small>F/Q/D</small>	almotriptan <small>F/Q/D</small> eletriptan <small>F/Q/D</small> Frova® <small>F/Q/D</small> frovatriptan <small>F/Q/D</small> Imitrex® <small>F/Q/D</small> Maxalt® <small>F/Q/D</small> Maxalt® MLT <small>F/Q/D</small> naratriptan <small>F/Q/D</small> Onzetra™ Xsail™ <small>F/Q/D</small> Relpax® sumatriptan-naproxen <small>F/Q/D</small>	<p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="background-color: #0070C0; color: white;">Agent</th><th style="background-color: #0070C0; color: white;">F/Q/D</th></tr> </thead> <tbody> <tr> <td>Onzetra™ Xsail™ 11 mg</td><td>16 units/30 days</td></tr> <tr> <td>almotriptan eletriptan (Relpax®) frovatriptan (Frova®) naratriptan rizatriptan (Maxalt®) rizatriptan (Maxalt® MLT) sumatriptan nasal spray (Imitrex®)</td><td>18 units/30 days</td></tr> </tbody> </table>	Agent	F/Q/D	Onzetra™ Xsail™ 11 mg	16 units/30 days	almotriptan eletriptan (Relpax®) frovatriptan (Frova®) naratriptan rizatriptan (Maxalt®) rizatriptan (Maxalt® MLT) sumatriptan nasal spray (Imitrex®)	18 units/30 days														
Agent	F/Q/D																					
Onzetra™ Xsail™ 11 mg	16 units/30 days																					
almotriptan eletriptan (Relpax®) frovatriptan (Frova®) naratriptan rizatriptan (Maxalt®) rizatriptan (Maxalt® MLT) sumatriptan nasal spray (Imitrex®)	18 units/30 days																					

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
	Tosymra™ F/Q/D Treximet® F/Q/D Zembrace™ SymTouch™ zolmitriptan F/Q/D Zomig® F/Q/D	sumatriptan (Imitrex®) sumatriptan-naproxen (Treximet®) Tosymra™ nasal spray zolmitriptan (Zomig®) Zomig® nasal spray
<b>Antipsychotics – Injectable</b>		
Abilify Maintena® Aristada® Aristada Initio® fluphenazine decanoate Haldol® decanoate haloperidol decanoate Invega Hafyera™ Invega Sustenna® Invega Trinza® Perseris™ Risperdal Consta® Zyprexa Relprevv®	Abilify Asimtufii® Uzedy™	
<b>Antipsychotics – Second Generation CC, ST</b>		
aripiprazole tablet DO asenapine (gen Saphris®) clozapine lurasidone (gen Latuda®) olanzapine tablet DO quetiapine F/Q/D quetiapine ER F/Q/D, DO risperidone ziprasidone capsule	Abilify® tablet DO Abilify MyCite® aripiprazole solution aripiprazole ODT Caplyta™ clozapine ODT Clozaril® Fanapt® Geodon® Invega® DO, F/Q/D Latuda® DO	<b>DOSE OPTIMIZATION (DO)</b> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected drugs and strengths</li> </ul> <b>CLINICAL CRITERIA (CC)</b> <ul style="list-style-type: none"> <li>Clinical editing will allow patients currently stabilized on a non-preferred agent to continue to receive that agent without PA</li> <li>Prior authorization is required when an oral SGA is utilized above the highest MDD according to FDA labeling.</li> <li>Prior authorization is required for patients less than 21 years of age when there is concurrent use of 2 or more different oral antipsychotics for greater than 90 days.</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters																																
<b>IV. Central Nervous System</b>																																		
	Lybalvi™ Nuplazid® olanzapine ODT <u>DO</u> paliperidone ER <u>F/Q/D, DO</u> Rexulti® <u>DO</u> Risperdal® Saphris® Secuado® <u>F/Q/D</u> Seroquel® <u>F/Q/D</u> Seroquel XR® <u>DO, F/q/d</u> Versacloz® Vraylar® <u>DO</u> Zyprexa® <u>DO</u> Zyprexa® Zydis	<ul style="list-style-type: none"> <li>Prior authorization is required for patients 21 years of age or older when 3 or more different oral second-generation antipsychotics are used for more than 180 days.</li> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> <li>PA is required for initial prescription for beneficiaries younger than the drug-specific minimum age as indicated below:</li> </ul> <table border="1" style="width: 100%; border-collapse: collapse;"> <tbody> <tr> <td style="padding: 2px;">aripiprazole (Abilify®)</td> <td style="padding: 2px; text-align: right;">6 years</td> </tr> <tr> <td style="padding: 2px;">aripiprazole (Abilify MyCite®)</td> <td style="padding: 2px; text-align: right;">18 years</td> </tr> <tr> <td style="padding: 2px;">asenapine (Saphris®)</td> <td style="padding: 2px; text-align: right;">10 years</td> </tr> <tr> <td style="padding: 2px;">asenapine (Secuado®)</td> <td style="padding: 2px; text-align: right;">18 years</td> </tr> <tr> <td style="padding: 2px;">brexpiprazole (Rexulti®)</td> <td style="padding: 2px; text-align: right;">13 years</td> </tr> <tr> <td style="padding: 2px;">cariprazine (Vraylar®)</td> <td style="padding: 2px; text-align: right;">18 years</td> </tr> <tr> <td style="padding: 2px;">clozapine (Clozaril®, Versacloz®)</td> <td style="padding: 2px; text-align: right;">12 years</td> </tr> <tr> <td style="padding: 2px;">ilooperidone (Fanapt®)</td> <td style="padding: 2px; text-align: right;">18 years</td> </tr> <tr> <td style="padding: 2px;">lumateperone (Caplyta™)</td> <td style="padding: 2px; text-align: right;">18 years</td> </tr> <tr> <td style="padding: 2px;">lurasidone HCl (Latuda®)</td> <td style="padding: 2px; text-align: right;">10 years</td> </tr> <tr> <td style="padding: 2px;">olanzapine (Zyprexa®)</td> <td style="padding: 2px; text-align: right;">10 years</td> </tr> <tr> <td style="padding: 2px;">paliperidone ER (Invega®)</td> <td style="padding: 2px; text-align: right;">12 years</td> </tr> <tr> <td style="padding: 2px;">pimavanserin (Nuplazid®)</td> <td style="padding: 2px; text-align: right;">18 years</td> </tr> <tr> <td style="padding: 2px;">quetiapine fum. (Seroquel®, Seroquel XR®)</td> <td style="padding: 2px; text-align: right;">10 years</td> </tr> <tr> <td style="padding: 2px;">risperidone (Risperdal®)</td> <td style="padding: 2px; text-align: right;">5 years</td> </tr> <tr> <td style="padding: 2px;">ziprasidone HCl (Geodon®)</td> <td style="padding: 2px; text-align: right;">10 years</td> </tr> </tbody> </table> <ul style="list-style-type: none"> <li>Require confirmation of diagnosis that supports the concurrent use of a Second Generation Antipsychotic and a CNS Stimulant for patients &lt; 18 years of age</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>For all Second Generation Antipsychotics used in the treatment of Major Depressive Disorder in the absence of other psychiatric comorbidities, trial with at least two different antidepressant agents is required</li> </ul>	aripiprazole (Abilify®)	6 years	aripiprazole (Abilify MyCite®)	18 years	asenapine (Saphris®)	10 years	asenapine (Secuado®)	18 years	brexpiprazole (Rexulti®)	13 years	cariprazine (Vraylar®)	18 years	clozapine (Clozaril®, Versacloz®)	12 years	ilooperidone (Fanapt®)	18 years	lumateperone (Caplyta™)	18 years	lurasidone HCl (Latuda®)	10 years	olanzapine (Zyprexa®)	10 years	paliperidone ER (Invega®)	12 years	pimavanserin (Nuplazid®)	18 years	quetiapine fum. (Seroquel®, Seroquel XR®)	10 years	risperidone (Risperdal®)	5 years	ziprasidone HCl (Geodon®)	10 years
aripiprazole (Abilify®)	6 years																																	
aripiprazole (Abilify MyCite®)	18 years																																	
asenapine (Saphris®)	10 years																																	
asenapine (Secuado®)	18 years																																	
brexpiprazole (Rexulti®)	13 years																																	
cariprazine (Vraylar®)	18 years																																	
clozapine (Clozaril®, Versacloz®)	12 years																																	
ilooperidone (Fanapt®)	18 years																																	
lumateperone (Caplyta™)	18 years																																	
lurasidone HCl (Latuda®)	10 years																																	
olanzapine (Zyprexa®)	10 years																																	
paliperidone ER (Invega®)	12 years																																	
pimavanserin (Nuplazid®)	18 years																																	
quetiapine fum. (Seroquel®, Seroquel XR®)	10 years																																	
risperidone (Risperdal®)	5 years																																	
ziprasidone HCl (Geodon®)	10 years																																	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
		<p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>• <b>asenapine (Secuado®)</b> 7.6 mg/24 hours</li> <li>• <b>lumateperone (Caplyta™)</b> 42 mg capsules: Maximum 1 unit/day</li> <li>• <b>paliperidone ER (Invega®)</b> 1.5 mg, 3 mg, 9 mg tablets: Maximum 1 unit/day</li> <li>• <b>paliperidone ER (Invega®)</b> 6 mg tablets: Maximum 2 units/day</li> <li>• <b>quetiapine/quetiapine ER (Seroquel®/Seroquel XR®)</b>: Minimum 100 mg/day; maximum 800 mg/day</li> <li>• <b>quetiapine (Seroquel®)</b>: Maximum 3 units per day, 90 units per 30 days</li> <li>• <b>quetiapine ER (Seroquel XR®)</b> 150 mg, 200 mg: 1 unit/day, 30 units/30 days</li> <li>• <b>quetiapine ER (Seroquel XR®)</b> 50 mg, 300 mg, 400 mg: 2 units/day, 60 units/30 days</li> </ul>
<b>Central Nervous System (CNS) Stimulants CC, F/Q/D</b>		
amphetamine salt combo IR (gen Adderall®) amphetamine salt combo ER (gen Adderall XR®) <b>DO</b> Concerta® <b>DO, BLTG</b> Daytrana® <b>BLTG</b> dexmethylphenidate (gen Focalin®) dexmethylphenidate ER <b>DO</b> (gen Focalin XR®) dextroamphetamine tablet methylphenidate solution (gen Methylin®) methylphenidate tablet (gen Ritalin®) methylphenidate ER (gen Aptensio® XR) Vyvanse® capsule, chewable <b>DO</b>	Adderall XR® <b>DO</b> Adzenys XR-ODT® amphetamine (gen Adzenys ER®) amphetamine (gen Evekeo®) Aptensio XR® armodafinil (gen Nuvigil®) Azstarys™ Cotempla® XR-ODT™ Desoxyn® Dexedrine® dextroamphetamine ER (gen Dexedrine®) dextroamphetamine solution (gen ProCentra®) dextroamphetamine tablet (gen Zenedi®) Dyanavel XR® Evekeo® Evekeo® ODT Focalin®	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved, compendia-supported and Medicaid covered indication</li> <li>• Prior authorization is required for initial prescriptions for stimulant therapy for beneficiaries <b>less than 3 years of age</b></li> <li>• Confirm diagnoses that support concurrent use of CNS Stimulant and Second Generation Antipsychotic agent for beneficiaries <b>less than 18 years of age</b></li> <li>• Patient-specific considerations for drug selection include treatment of excessive sleepiness associated with shift work sleep disorder, narcolepsy, or as an adjunct to standard treatment for obstructive sleep apnea.</li> <li>• PA required for initiation of CNS Stimulant for patients currently on an opioid</li> <li>• PA required for initiation of CNS Stimulant for patients currently on a benzodiazepine</li> </ul> <p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>• See Dose Optimization Chart for affected drugs and strengths</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
	Focalin XR® <b>DO</b> Jornay PM™ methamphetamine (gen Desoxyn®) Methyltin® methylphenidate (gen Daytrana®) methylphenidate chewable tablet (gen Methyltin®) methylphenidate CD <b>DO</b> methylphenidate ER 45 mg, 63 mg, 72 mg tablet methylphenidate ER (gen Concerta®, Ritalin LA®, Metadate®) modafinil (gen Provigil®) <b>DO</b> Mydayis™ Nuvigil® ProCentra® Provigil® <b>DO</b> QuilliChew ER™ <b>DO</b> Quillivant XR® Relexxi® Ritalin® Ritalin LA® <b>DO</b> Sonusi™ Wakix® Xelstrym™ Zenzedi®	<b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b> <ul style="list-style-type: none"> <li>Quantity limits based on daily dosage as determined by FDA labeling</li> <li>Quantity limits to include: <ul style="list-style-type: none"> <li>Short-acting CNS stimulants: not to exceed 3 dosage units daily with maximum of 90 days per strength (for titration)</li> <li>Long-acting CNS stimulants: not to exceed 1 dosage unit daily with maximum of 90 days. Concerta 36mg and Cotempla XR-ODT 25.9 mg.; not to exceed 2 units daily</li> <li>Azstarys; not to exceed 1 dosage unit per day</li> <li>Pitolisant (Wakix®): not to exceed 2 dosage units daily of the 17.8 mg tablets or 3 dosage units daily of the 4.45 mg tablets.</li> </ul> </li> </ul>
<b>Movement Disorder Agents <b>CC</b></b>		
Austedo® Ingrezza® Ingrezza® titration pack tetrabenazine	Austedo® XR Xenazine®	<b>CLINICAL CRITERIA (CC)</b> <ul style="list-style-type: none"> <li>Confirm diagnosis for an FDA-approved or compendia-supported indication</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
<b>Multiple Sclerosis Agents</b>		
Avonex® Betaseron® Copaxone® 20 mg/mL <sup>BLTG</sup> dimethyl fumarate DR	Aubagio® Bafiertam™ Copaxone® 40 mg/mL Extavia® fingolimod (gen Gilenya®) Gilenya® glatiramer Kesimpta® Mavenclad® Mayzent® Plegridy® Ponvory™ F/Q/D Rebif® Rebif® Rebidose® Tascenso ODT™ Tecfidera® teriflunomide (gen Aubagio®) Vumerity® Zeposia® CC, ST	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>• <b>Zeposia® (ozanimod)</b>: Confirm diagnosis for FDA- or compendia-supported use</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>• <b>Zeposia® (ozanimod)</b>: For an indication of Ulcerative Colitis             <ul style="list-style-type: none"> <li>• Trial of a non-specific anti-inflammatory drug such as an aminosalicylate or immunosuppressant, or a disease-modifying anti-rheumatic drug (DMARD), and;</li> <li>• Trial of a preferred systemic immunomodulator</li> </ul> </li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>• <b>Ponvory™ (ponesimod) starter pack</b>; maximum quantity is 14, no refills</li> <li>• <b>Ponvory™ (ponesimod)</b>; maintenance limited to a 30-day supply</li> </ul>
<b>Non-Ergot Dopamine Receptor Agonists</b>		
pramipexole ropinirole	Kynmobi™ CC Mirapex ER® Neupro® pramipexole ER ropinirole ER	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>• <b>apomorphine (Kynmobi™)</b>: Confirm diagnosis of FDA-approved, compendia-supported, and Medicaid-covered indication</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
<b>Other Agents for Attention Deficit Hyperactivity Disorder (ADHD) <small>CC</small></b>		
atomoxetine <small>DO</small> clonidine ER guanfacine ER <small>DO</small>	Intuniv® <small>DO</small> Qelbree™ Strattera® <small>DO</small>	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Confirm diagnosis for an FDA-approved or compendia-supported indication for beneficiaries &lt; 18 years of age.</li> <li>Prior authorization is required for initial prescriptions for non-stimulant therapy for beneficiaries <b>less than 6 years of age</b></li> </ul> <p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul>
<b>Sedative Hypnotics/Sleep Agents <small>F/Q/D</small></b>		
estazolam <small>CC</small> temazepam 15 mg, 30 mg <small>CC</small> zolpidem tablet <small>CC</small>	Ambien® <small>CC</small> Ambien CR® <small>CC</small> Belsomra® Dayvigo™ Doral® <small>CC</small> doxepin (gen Silenor®) Edluar® <small>CC</small> eszopiclone Halcion® <small>CC</small> Lunesta® <small>DO</small> quazepam <small>CC</small> (gen Doral®) Quviquiq™ ramelteon (gen Rozerem®) Restoril® <small>CC</small> Rozerem® Silenor® temazepam 7.5 mg, 22.5 mg <small>CC</small> triazolam <small>CC</small> zaleplon zolpidem sublingual, capsule <small>CC</small> zolpidem ER <small>CC</small>	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul> <p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li><b>Zolpidem products:</b> Confirm dosage is consistent with FDA labeling for initial prescriptions</li> <li><b>Benzodiazepine Agents</b> (estazolam, Halcion®, Restoril®, temazepam, triazolam): <ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> <li>PA required for initiation of benzodiazepine therapy in patients currently on opioid or oral buprenorphine therapy</li> <li>PA required for any additional benzodiazepine prescription in patients currently on benzodiazepine therapy</li> <li>PA required when greater than a 14-day supply of a benzodiazepine is prescribed for someone on a CNS stimulant</li> </ul> </li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>Frequency and duration limits for the following products: <ul style="list-style-type: none"> <li>For <b>non-zaleplon</b> and <b>non-benzodiazepine</b> containing products: <ul style="list-style-type: none"> <li>30 dosage units per fill/1 dosage unit per day/30 days</li> </ul> </li> <li>For <b>zaleplon</b>-containing products: <ul style="list-style-type: none"> <li>60 dosage units per fill/2 dosage units per day/30 days</li> </ul> </li> </ul> </li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
		<ul style="list-style-type: none"> <li>– Duration limit equivalent to the maximum recommended duration:           <ul style="list-style-type: none"> <li>o 180 days for immediate-release <b>zolpidem</b> (Ambien®, Edluar®) products</li> <li>o 180 days for <b>eszopiclone</b> and <b>ramelteon</b> (Rozerem®) products</li> <li>o 180 days for lemborexant (Dayvigo™)</li> <li>o 168 days for <b>zolpidem ER</b> (Ambien CR®) products</li> <li>o 90 days for daridorexant (Quviquiq™)</li> <li>o 90 days for <b>suvorexant</b> (Belsomra®)</li> <li>o 90 days for <b>doxepin</b> (Silenor®)</li> <li>o 30 days for zaleplon (Sonata®) products</li> <li>o 30 days for benzodiazepine agents (estazolam, Halcion®, Restoril®, temazepam, triazolam) for the treatment of insomnia</li> </ul> </li> </ul> <p>Additional/Alternate parameters:</p> <ul style="list-style-type: none"> <li>• For patients naïve to non-benzodiazepine sedative hypnotics (NBSH): First-fill duration and quantity limit of 10 dosage units as a 10-day supply, except for zaleplon-containing products which the quantity limit is 20 dosage units as a 10-day supply</li> </ul>
<b>Selective Serotonin Reuptake Inhibitors (SSRIs)</b>		
citalopram tablet, solution escitalopram tablet fluoxetine capsule, solution paroxetine tablet sertraline tablet, concentrate Viibryd® <small>DO, BLTG, 1</small>	Celexa® citalopram capsule escitalopram solution fluoxetine tablet fluoxetine DR weekly fluvoxamine <small>CC</small> fluvoxamine ER <small>CC</small> Lexapro® <small>DO</small> paroxetine capsule paroxetine CR paroxetine suspension Paxil® Paxil CR® Pexeva®	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>• See Dose Optimization Chart for affected strengths</li> </ul> <p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>• Clinical editing will allow patients currently stabilized on fluvoxamine or fluvoxamine ER to continue to receive that agent without PA</li> <li>• Clinical editing to allow patients with a diagnosis of Obsessive-Compulsive Disorder (OCD) to receive fluvoxamine and fluvoxamine ER without prior authorization</li> </ul>

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf)

26

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IV. Central Nervous System</b>		
	Prozac® sertraline capsule Trintellix® <small>DO</small> vilazodone (gen Viibryd®) Zoloft®	
<b>Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs)</b>		
duloxetine 20 mg, 30 mg, 60 mg (gen Cymbalta®) venlafaxine venlafaxine ER capsule	Cymbalta® desvenlafaxine ER desvenlafaxine succinate ER <small>DO</small> Drizalma Sprinkle™ duloxetine 40 mg Effexor XR® <small>DO</small> Fetzima® Pristiq® <small>DO</small> Savella® venlafaxine ER tablet	<b>DOSE OPTIMIZATION (DO)</b> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>V. Dermatologic Agents</b>		
<b>Acne Agents, Topical</b>		
adapalene/benzoyl peroxide adapalene cream adapalene OTC gel Retin-A® cream <small>CC, BLTG</small> tazarotene cream <small>CC</small> tretinooin gel (gen Avita, Retin-A) <small>CC</small>	adapalene Rx gel, gel pump adapalene/benzoyl peroxide Altreno® <small>CC</small> Arazlo™ <small>CC</small> Atralin® <small>CC</small> Avita® <small>CC</small> clindamycin/tretinooin <small>CC</small> dapson Fabior® <small>CC</small> Retin-A® gel <small>CC</small> Retin-A Micro® <small>CC</small> tazarotene foam (gen Fabior®) <small>CC</small> tazarotene gel <small>CC</small> tretinooin cream, gel <small>CC</small> (gen Atralin) tretinooin micro <small>CC</small> Winlevi® Ziana® <small>CC</small>	<b>CLINICAL CRITERIA</b> <ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved, compendia-supported, and Medicaid-covered indication</li> </ul>
<b>Actinic Keratosis Agents</b>		
diclofenac 3% gel fluorouracil solution fluorouracil 0.5% cream (gen Carac) fluorouracil 5% cream (gen Efudex cream) imiquimod (gen Aldara)	Carac® Efudex® imiquimod (gen Zyclara) Tolak® Zyclara®	
<b>Antibiotics – Topical</b>		
mupirocin ointment	Centany® mupirocin cream Xepi™	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>V. Dermatologic Agents</b>		
<b>Anti-Fungals – Topical</b>		
ciclopirox cream, suspension, shampoo clotrimazole OTC clotrimazole Rx clotrimazole/betamethasone cream ketoconazole cream ketoconazole 2% shampoo miconazole OTC nystatin cream, ointment, powder nystatin/triamcinolone terbinafine OTC tolnaftate OTC	Alevazol OTC butenafine (gen Mentax®) Cyclodan® cream ciclopirox gel clotrimazole/betamethasone lotion econazole Ertaczo® Extina® ketoconazole foam Loprox® shampoo luliconazole Luzu® Mentax® miconazole/zinc/petrolatum (gen Vusion®) F/Q/D naftifine Naftin® oxiconazole Oxistat® Vusion® F/Q/D	<b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b> • <b>Vusion® 50 gm ointment</b> –Maximum 100 grams in a 90-day time period

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>V. Dermatologic Agents</b>		
<b>Anti-Infectives – Topical</b>		
clindamycin solution, gel, lotion, swab clindamycin/benzoyl peroxide (gen Duac®) erythromycin solution, gel	Acanya® Benzamycin® Cleocin T® clindamycin foam clindamycin/benzoyl peroxide (gen BenzaClin®) clindamycin/benzoyl peroxide (gen Acanya®) Erygel® erythromycin swab erythromycin/benzoyl peroxide Evoclin® Neuac® Onexton®	
<b>Anti-Virals – Topical</b>		
acyclovir cream docosanol (gen Abreva)	acyclovir ointment Denavir® penciclovir (gen Denavir®) Sitavig® Xerese® Zovirax® cream, ointment	
<b>Immunomodulators – Topical <span style="color: red;">CC</span></b>		
pimecrolimus tacrolimus	Elidel® Protopic®	<b>CLINICAL CRITERIA</b> <ul style="list-style-type: none"> <li>• All prescriptions require prior authorization</li> <li>• Refills on prescriptions are allowed</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>V. Dermatologic Agents</b>		
<b>Psoriasis Agents – Topical</b>		
calcipotriene cream, ointment, scalp solution	calcipotriene foam (gen Sorilux®) calcipotriene/betamethasone dipropionate (gen Taclonex®) calcitriol ointment (gen Vectical®) Dovonex® Duobrii™ Enstilar® Sorilux® Taclonex® Vtama® Zoryve™	
<b>Steroids, Topical – Low Potency</b>		
hydrocortisone acetate OTC hydrocortisone acetate Rx	alclometasone Derma-Smoothe/FS® desonide fluocinolone oil Texacort®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>V. Dermatologic Agents</b>		
<b>Steroids, Topical – Medium Potency</b>		
mometasone furoate	Beser lotion betamethasone valerate foam clocortolone Cloderm® fluocinolone acetonide cream, ointment, solution flurandrenolide fluticasone propionate hydrocortisone butyrate cream, lotion, ointment, solution hydrocortisone valerate Locoid® Locoid Lipocream® Luxiq® Pandel® prednicarbate Synalar®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>V. Dermatologic Agents</b>		
<b>Steroids, Topical – High Potency</b>		
betamethasone dipropionate lotion, cream betamethasone dipropionate augmented cream betamethasone valerate cream, ointment triamcinolone acetonide	amcinonide ApexiCon-E® betamethasone dipropionate gel, ointment betamethasone dipropionate augmented lotion, ointment betamethasone valerate lotion desoximetasone diflorasone Diprolene® fluocinonide 0.1% cream (gen Vanos®) fluocinonide ointment, cream, gel, solution, emollient halcinonide cream (gen Halog®) Halog® cream, solution, ointment Kenalog® Topicort® triamcinolone spray Vanos®	
<b>Steroids, Topical – Very High Potency</b>		
clobetasol cream, emollient, gel, ointment, solution halobetasol cream, ointment	Bryhali™ clobetasol foam, lotion, spray, shampoo Clobex® halobetasol foam Impeklo™ Lexette™ foam Olux® Olux-E® Temovate® Ultravate®	

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VI. Endocrine and Metabolic Agents</b>		
<b>Anabolic Steroids – Topical <small>CDRP, F/Q/D</small></b>		
testosterone gel testosterone pump	Androderm® AndroGel® pump Fortesta® Testim® Vogelxo	<p><b>CLINICAL DRUG REVIEW PROGRAM (CDRP)</b></p> <ul style="list-style-type: none"> <li>For diagnosis of hypogonadotropic or primary hypogonadism:             <ul style="list-style-type: none"> <li>Requires documented low testosterone concentration with two tests prior to initiation of therapy.</li> <li>Require documented testosterone therapeutic concentration to confirm response after initiation of therapy.</li> </ul> </li> <li>For diagnosis of delayed puberty:             <ul style="list-style-type: none"> <li>Requires documentation that growth hormone deficiency has been ruled out prior to initiation of therapy.</li> <li>1.62% gel only: For diagnosis of gender dysphoria please refer to <a href="#">July 2020 edition of the Medicaid Update</a>: <a href="https://www.health.ny.gov/health_care/medicaid/program/update/2020/no12_2020-07.htm#transgender">https://www.health.ny.gov/health_care/medicaid/program/update/2020/no12_2020-07.htm#transgender</a></li> </ul> </li> </ul> <p>The Anabolic Steroid fax form can be found at:  <a href="https://newyork.fhsc.com/downloads/providers/NYRx_CDRP_PA_Worksheet_Prescribers_Anabolic_Steroids.pdf">https://newyork.fhsc.com/downloads/providers/NYRx_CDRP_PA_Worksheet_Prescribers_Anabolic_Steroids.pdf</a></p> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>Limitations for anabolic steroid products based on approved FDA labeled daily dosing and documented diagnosis:             <ul style="list-style-type: none"> <li>Duration limit of 6 months for delayed puberty</li> </ul> </li> </ul>
<b>Biguanides</b>		
Glumetza® <small>BLTG</small> metformin HCl metformin ER (gen Glucophage XR®)	metformin solution (gen Riomet®) metformin ER <small>DO</small> (gen Fortamet®, Glumetza®) Riomet® Riomet ER™	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VI. Endocrine and Metabolic Agents</b>		
<b>Bisphosphonates – Oral</b>		
alendronate	Actonel® Atelvia® Boniva® Fosamax® Fosamax® Plus D ibandronate risedronate	
<b>Dipeptidyl Peptidase-4 (DPP-4) Inhibitors <sup>ST</sup></b>		
Glyxambi® Janumet® Janumet® XR Januvia® <sup>DO</sup> Jentadueto® Jentadueto® XR <sup>1</sup> Kazano® <sup>BLTG</sup> Kombiglyze® XR <sup>1</sup> Nesina® <sup>BLTG</sup> Onglyza® <sup>DO, 1</sup> Oseni® <sup>BLTG</sup> Tradjenta®	alogliptin alogliptin/metformin alogliptin/pioglitazone Qtern® Steglujan®	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>Requires a trial with metformin with or without insulin prior to DPP-4 Inhibitor therapy unless there is a documented contraindication.</li> </ul>

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf)

35

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VI. Endocrine and Metabolic Agents</b>		
<b>Glucagon Agents</b>		
Baqsimi® <sup>1</sup> glucagon vial glucagon HCl emergency kit (Fresenius) Gvoke® <sup>1</sup> pen, syringe, vial Zeg掬ogue® pen, syringe	glucagon emergency kit (Eli Lilly, Amphastar)	
<b>Glucagon-like Peptide-1 (GLP-1) Agonists CC, ST</b>		
Byetta® Ozempic® Trulicity® Victoza®	Adlyxin® Bydureon® BCise™ Mounjaro® Rybelsus® Soliqua® Xultophy®	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved, compendia-supported, and Medicaid-covered indication</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>Trial with metformin with or without insulin prior to initiating GLP-1 agonist therapy unless there is a contraindication, or the drug is being used for an FDA-approved Medicaid covered indication other than, or in addition to, Type 2 Diabetes.</li> </ul>

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf) 36

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VI. Endocrine and Metabolic Agents</b>		
<b>Glucocorticoids – Oral</b>		
budesonide EC, DR dexamethasone tablet hydrocortisone methylprednisolone dose-pack prednisolone solution prednisone dose-pack, tablet	Alkindi® Sprinkle budesonide ER Cortef® cortisone dexamethasone elixir, solution dexamethasone intensol Emflaza® Hemady™ Medrol® dose-pack, tablet methylprednisolone 4 mg, 8 mg, 16 mg, 32 mg Millipred® Millipred® DP Ortikos™ prednisolone ODT prednisolone tablet (gen Millipred®) prednisone intensol, solution Rayos® Uceris®	
<b>Growth Hormones CC, CDRP</b>		
Genotropin® Norditropin®	Humatropे® Nutropin AQ® Omnitrope® Saizen® Skytrofa® Sogroya® Zomacton®	<p><b>CLINICAL DRUG REVIEW PROGRAM (CDRP)</b></p> <ul style="list-style-type: none"> <li>Prescribers or their authorized agents may call or submit a fax request for a PA for beneficiaries 18 years of age or older</li> </ul> <p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Patient-specific considerations for drug selection include concerns related to use of a non-preferred agent for FDA-approved indications that are not listed for a preferred agent.</li> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VI. Endocrine and Metabolic Agents</b>		
<b>Insulin – Long-Acting</b>		
insulin glargine solostar, vial (gen Lantus® Solostar®, vial) Lantus® Solostar®, vial Levemir®	Basaglar® insulin degludec vial, pen (gen Tresiba) insulin glargine-YFGN: vial, pen Rezvoglar™ Semglee®-YFGN: vial, pen Toujeo® Solostar® Toujeo® Max Solostar® Tresiba®	
<b>Insulin – Mixes</b>		
Humalog® 50/50 Mix: pen and vial Humalog® 75/25 Mix: vial insulin lispro 75/25 mix: pen (gen Humalog® Mix) insulin aspart prot/insulin aspart: vial, pen (gen Novolog)	Humalog® 75/25 mix: pen Novolog® Mix: vial, pen	
<b>Insulin – Rapid-Acting</b>		
Apidra® Humalog® Jr. 100 U/mL Kwikpen Humalog® 100 U/mL vial, pen, cartridge insulin aspart (gen Novolog®) cartridge, vial, pen insulin lispro (gen Humalog® U100) vial, pen insulin lispro junior (gen Humalog® Jr.) Novolog® cartridge, vial, FlexPen	Admelog® Afrezza® Fiasp® Penfill, FlexTouch Humalog® 200 U/mL Lyumjev™	
<b>Pancreatic Enzymes</b>		
Creon® Zenpep®	Pertzye® Viokace®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VI. Endocrine and Metabolic Agents</b>		
<b>Sodium Glucose Co-Transporter 2 (SGLT2) Inhibitors <sup>ST</sup></b>		
Farxiga® Invokana® Jardiance®	Inpefa™ Invokamet® Invokamet® XR Segluromet® Steglatro® Synjardy® Synjardy® XR Trijardy® XR Xigduo® XR	<p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>Requires trial with metformin with or without insulin prior to initiating SGLT2 therapy unless there is a contraindication, or the drug is being used for an FDA-approved Medicaid covered indication other than, or in addition to, Type 2 Diabetes.</li> </ul>
<b>Thiazolidinediones (TZDs) <sup>ST</sup></b>		
pioglitazone	ACTOplus Met® Actos® <sup>DO</sup> Duetact® pioglitazone/glimepiride pioglitazone/metformin	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>Requires a trial with metformin with or without insulin prior to initiating TZD therapy unless there is a documented contraindication.</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VII. Gastrointestinal</b>		
<b>Anti-Emetics</b>		
aprepitant pack Diclegis® <small>CC</small> , doxylamine succ/pyridoxine ondansetron ODT, solution, tablet	Akynzeo® Anzemet® aprepitant capsule Bonjesta® <small>CC</small> Emend® capsule, powder packet, TriPack granisetron tablet Sancuso®	<b>CLINICAL CRITERIA (CC)</b> <ul style="list-style-type: none"> <li>• <b>Diclegis® and Bonjesta®:</b> Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul>
<b>Gastrointestinal Antibiotics</b>		
Firvanq® <small>BLTG</small> metronidazole tablet neomycin vancomycin capsule	Difidid® Flagyl® metronidazole capsule nitazoxanide paromomycin tinidazole Vancocin® vancomycin solution Xifaxan® <small>CC, ST, F/Q/D</small>	<b>CLINICAL CRITERIA (CC)</b> <ul style="list-style-type: none"> <li>• <b>Xifaxan®:</b> Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul> <b>STEP THERAPY (ST)</b> <ul style="list-style-type: none"> <li>• <b>Xifaxan®:</b> Requires trial of a fluoroquinolone antibiotic or azithromycin before Xifaxan® for treatment of Traveler's Diarrhea</li> </ul> <b>QUANTITY LIMITS:</b> <p><b>Xifaxan®:</b></p> <ul style="list-style-type: none"> <li>• Irritable bowel syndrome with diarrhea (550 mg tablets) – 42 tablets per 30 days (Dose = 550 mg three times a day for 14 days) <ul style="list-style-type: none"> <li>– Maximum of 42 days' supply (126 units) per 365 days (3 rounds of therapy).</li> </ul> </li> <li>• Small Intestine Bacterial Overgrowth (550mg tablets) - 42 tablets per 30 days (Dose = 550mg three times a day for 10-14 days); <ul style="list-style-type: none"> <li>– Maximum of 28 days' supply (84 units) per 365 days (2 rounds of therapy).</li> </ul> </li> </ul>
<b>Helicobacter pylori Agents</b>		
Pylera® <small>BLTG</small>	bismuth/metronidazole/tetracycline (gen Pylera®) lansoprazole/amoxicillin/clarithromycin Omeclamox-Pak® Talicia®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VII. Gastrointestinal</b>		
<b>Proton Pump Inhibitors (PPIs) F/Q/D</b>		
Dexilant® <small>DO, BLTG, 1</small> esomeprazole magnesium Rx capsule <sup>1</sup> lansoprazole Rx capsule <sup>1</sup> omeprazole Rx pantoprazole tablet Protonix suspension <small>BLTG</small> rabeprazole <sup>1</sup> Zegerid® Rx <small>BLTG</small>	Aciphex® dexlansoprazole (gen Dexilant) esomeprazole magnesium tablet OTC esomeprazole capsules OTC esomeprazole suspension Konvomep™ lansoprazole Rx ODT Nexium® RX <small>DO</small> omeprazole OTC omeprazole/sodium bicarbonate Rx omeprazole/sodium bicarbonate OTC pantoprazole suspension Prevacid® OTC Prevacid® Rx <small>DO</small> Prilosec® Rx Protonix® tablet	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>Quantity limits: <ul style="list-style-type: none"> <li>Once daily dosing for: <ul style="list-style-type: none"> <li>GERD</li> <li>erosive esophagitis</li> <li>healing and maintenance of duodenal/gastric ulcers (including NSAID-induced)</li> <li>prevention of NSAID-induced ulcers</li> </ul> </li> <li>Twice daily dosing for: <ul style="list-style-type: none"> <li>hypersecretory conditions</li> <li>Barrett's esophagitis</li> <li>H. pylori</li> <li>refractory GERD</li> </ul> </li> </ul> </li> <li>Duration limits: <ul style="list-style-type: none"> <li>90 days for: <ul style="list-style-type: none"> <li>GERD</li> </ul> </li> <li>365 days for: <ul style="list-style-type: none"> <li>Maintenance treatment of duodenal ulcers, or erosive esophagitis</li> </ul> </li> <li>14 days for: <ul style="list-style-type: none"> <li>H. pylori</li> </ul> </li> </ul> </li> </ul>

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf) 41

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VII. Gastrointestinal</b>		
<b>Sulfasalazine Derivatives</b>		
Apriso® <a href="#">BLTG</a> Lialda® <a href="#">BLTG</a> Pentasa® <a href="#">BLTG</a> sulfasalazine DR sulfasalazine IR	Asacol HD® Azulfidine® Azulfidine Entab® balsalazide Colazal® Delzicol® Dipentum® mesalamine DR (gen Delzicol®) mesalamine DR (gen Lialda®) mesalamine ER (gen Apriso®) mesalamine ER (gen Pentasa®) mesalamine DR	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VIII. Hematological Agents</b>		
<b>Anticoagulants – Injectable F/Q/D</b>		
enoxaparin sodium Fragmin® vial	Arixtra® <small>CC</small> fondaparinux <small>CC</small> Fragmin® syringe Lovenox®	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>For patients requiring &gt; 30 days of therapy: Require confirmation of FDA-approved or compendia-supported indication</li> <li><b>Arixtra® (fondaparinux)</b> Clinical editing to allow patients with a diagnosis of Heparin Induced Thrombocytopenia (HIT) to receive therapy without prior authorization.</li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>Duration Limit: No more than 30 days for members initiating therapy</li> </ul>
<b>Anticoagulants – Oral</b>		
Eliquis® Pradaxa® capsule <small>BLTG</small> warfarin Xarelto® tablet <small>DO</small>	dabigatran (gen Pradaxa®) Pradaxa® pellet pack Savaysa® Xarelto® dose pack, suspension	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul>
<b>Colony Stimulating Factors</b>		
Neupogen® Nyvepria™	Fylnetra® Fulphila™ Granix® Leukine® Neulasta® Nivestym™ Releuko™ Stimufend® Udenyca® Zarxio® Ziextenzo®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VIII. Hematological Agents</b>		
<b>Erythropoiesis Stimulating Agents (ESAs) <sup>CC</sup></b>		
Aranesp® <sup>1</sup> Epogen® Retacrit®	Mircera® Procrit®	<b>CLINICAL CRITERIA (CC)</b> <ul style="list-style-type: none"> <li>Confirm diagnosis for FDA- or compendia-supported uses</li> </ul>
<b>Hemophilia Agents – Factor VIII</b>		
Advate® Adynovate® Afystyla® Eloctate® Esperoct® Hemofil® M Humate-P® Jivi® Koate® Kogenate® FS Kovaltry® Novoeight® Nuwiq® Obizur® Recombinate™ Xyntha® Xyntha® Solofuse	Altuviiio™	

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>VIII. Hematological Agents</b>		
<b>Hemophilia Agents – Factor IX</b>		
AlphaNine® SD Alprolix® BeneFIX® Idelvion® Ixinity® Profilnine® Rebinyn® Rixubis®	N/A	
<b>Hemophilia Agents – Other</b>		
Alphanate® (von Willebrand factor/Factor VIII) Coagadex® (Factor X) Corifact® (Factor XIII) Feiba® NF (activated prothrombin complex) Hemlibra® (emicizumab-kxwh) Novoseven® RT (Factor VIIa) Sevenfact® (Factor VIIa-jncw) Tretten® (Factor XIII) Vonvendi® (von Willebrand factor) Wilate® (von Willebrand factor/Factor VIII)	N/A	
<b>Platelet Inhibitors</b>		
Brilinta® clopidogrel dipyridamole dipyridamole/aspirin	Effient® Plavix® prasugrel	

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf) 45

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IX. Immunologic Agents</b>		
<b>Immunomodulators – Systemic <small>CC, ST</small></b>		
Cosentyx® Dupixent® Enbrel® Fasenra® Humira® Nucala® Xolair®	Actemra® subcutaneous adalimumab-FKJP adalimumab-ADAZ Adbry™ Amjevita™ Cibinquo™ Cimzia® Cyltezo® Hadlima™ Hulio® (adalimumab-FKJP) Hyrimoz® (adalimumab-ADAZ) Idacio® Yuflyma® Yusimry™ Illumya® Kevzara® Kineret® Olumiant® Orencia® subcutaneous Otezla® Rinvoq™ ER Siliq™ Siponimod® Skyrizi® Skyrizi® On-Body Sotyktu™ Stelara® Taltz® Tezspire® pen Tremfya®	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Confirm diagnosis for FDA- or compendia-supported uses</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <p>For indications not specified below</p> <ul style="list-style-type: none"> <li>Trial of a non-specific anti-inflammatory drug such as an aminosalicylate or immunosuppressant, or a disease-modifying anti-rheumatic drug (DMARD)</li> <li>Trial of a TNF inhibitor prior to treatment with a JAK inhibitor</li> </ul> <p><b>INDICATION-SPECIFIC REQUIREMENTS:</b></p> <ul style="list-style-type: none"> <li>Asthma: <ul style="list-style-type: none"> <li>history and concurrent use of a corticosteroid</li> </ul> </li> <li>Nasal polyps: <ul style="list-style-type: none"> <li>history and concurrent use of an intranasal corticosteroid</li> </ul> </li> <li>Atopic dermatitis: <ul style="list-style-type: none"> <li>Trial with a topical prescription product for a duration of at least 3 months.</li> <li>For JAK inhibitors: Trial of topical prescription product and systemic product for a combined duration of at least 6 months.</li> </ul> </li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>IX. Immunologic Agents</b>		
<b>Immunomodulators – Systemic CC, ST</b>		
	Xeljanz® Xeljanz® XR	
<b>Immunosuppressives, Oral</b>		
azathioprine CellCept® suspension <small>BLTG</small> cyclosporine softgel, capsule cyclosporine modified capsule, solution mycophenolic acid <sup>1</sup> mycophenolate mofetil capsule, tablet Rapamune® solution <small>BLTG</small> Rapamune® tablet <sup>1</sup> sirolimus tablet tacrolimus	Astagraf XL® Azasan® CellCept® capsule, tablet Envarsus XR® everolimus (gen Zortress®) Imuran® Lupkynis™ <small>CC, ST, F/Q/D</small> mycophenolate mofetil suspension Myfortic® Neoral® Prograf® Sandimmune® solution, capsule sirolimus solution Zortress®	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Lupkynis™ (voclosporin) – Confirm diagnosis for FDA- or compendia-supported uses</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>Trial of mycophenolate prior to Lupkynis™</li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>Lupkynis™ limited to 30-day supply</li> </ul>

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf) 47

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>X. Miscellaneous Agents</b>		
<b>Progestins (for Cachexia)</b>		
megestrol acetate suspension	megestrol 625 mg/5 mL suspension	
<b>Epinephrine – Self-injected</b>		
EpiPen® <small>BLTG</small> EpiPen Jr.® <small>BLTG</small>	Auvi-Q® (0.1mg) epinephrine (gen Adrenaclick®) epinephrine (gen EpiPen®) epinephrine (gen EpiPen Jr.®) Symjepi®	

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf) 48

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XI. Musculoskeletal Agents</b>		
<b>Skeletal Muscle Relaxants</b>		
baclofen tablet chlorzoxazone 500 mg cyclobenzaprine 5 mg, 10 mg tablet dantrolene methocarbamol orphenadrine ER tizanidine tablet	Amrix® baclofen solution F/Q/D baclofen (gen Fleqsuvy™) carisoprodol ST, F/Q/D carisoprodol compound ST, F/Q/D carisoprodol compound/codeine CC, ST, F/Q/D  chlorzoxazone (gen Lorzone) 375 mg, 750 mg cyclobenzaprine 7.5 mg cyclobenzaprine ER capsule (gen Amrix) Dantrium® Fexmid® Fleqsuvy™ Lorzone® Lyvispah™ metaxalone orphenadrine-aspirin-caffeine Soma® ST, F/Q/D Soma® 250 ST, F/Q/D tizanidine capsule Zanaflex®	<p><b>CLINICAL CRITERIA (CC)</b></p> <p>For carisoprodol/codeine products:</p> <ul style="list-style-type: none"> <li>• Limited to a total of 4 opioid prescriptions every 30 days; exemption for diagnosis of cancer or sickle cell disease</li> <li>• Medical necessity rationale for opioid therapy is required for patients on established opioid dependence therapy</li> <li>• PA required for initiation of opioid therapy in patients currently on benzodiazepine therapy</li> <li>• PA required for any codeine containing products in patients &lt; 12 years</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>• Trial with 1 preferred analgesic and 2 preferred skeletal muscle relaxants prior to use of <b>carisoprodol</b> containing products: <ul style="list-style-type: none"> <li>– carisoprodol</li> <li>– carisoprodol/ASA</li> <li>– carisoprodol/ASA/codeine</li> <li>– Soma®</li> </ul> </li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>• Maximum 84 cumulative units per a year</li> <li>• Baclofen solution – Maximum 946 mL per 30 days</li> <li>• <b>Carisoprodol</b> – Maximum 4 units per day, 21-day supply</li> <li>• <b>Carisoprodol combinations</b> – Maximum 8 units per day, 21-day supply (not to exceed the 84 cumulative units per year limit)</li> </ul>

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XII. Ophthalmics</b>		
<b>Alpha-2 Adrenergic Agonists (for Glaucoma) – Ophthalmic</b>		
Alphagan P® 0.1% Alphagan P® 0.15% <small>BLTG</small> brimonidine 0.2% Simbrinza®	apraclonidine brimonidine P 0.15% lopidine®	
<b>Antibiotics – Ophthalmic</b>		
bacitracin/polymyxin B erythromycin gentamicin Natacyn® neomycin/gramicidin/polymyxin polymyxin(trimethoprim sulfacetamide solution tobramycin	Azasite® bacitracin neomycin/bacitracin/polymyxin Polytrim® sulfacetamide ointment Tobrex®	
<b>Antibiotics/Steroid Combinations – Ophthalmic</b>		
Blephamide® neomycin/polymyxin/dexamethasone sulfacetamide/prednisolone TobraDex® ointment tobramycin/dexamethasone suspension	Maxitrol® neomycin / bacitracin/polymyxin /HC neomycin/polymyxin/HC Pred-G® TobraDex® ST TobraDex® suspension Zylet®	

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XII. Ophthalmics</b>		
<b>Antihistamines – Ophthalmic</b>		
azelastine <sup>1</sup> ketotifen OTC <sup>1</sup> olopatadine OTC	bepotastine (gen Bepreve®) Bepreve® epinastine Lastacift® olopatadine Rx Pataday® Zaditor® OTC Zerviate™	
<b>Anti-inflammatories/Immunomodulators – Ophthalmic CC, F/Q/D</b>		
Restasis® <sup>BLTG</sup> Restasis MultiDose® Xiidra®	Cequa® cyclosporine (gen Restasis®) Tyrvaya™ Verkazia®	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>Diagnosis documentation required to justify utilization as a first line agent or attempt treatment with an artificial tear, gel, or ointment.</li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>Cequa®, Restasis®, Xiidra®: 60 vials dispensed as a 30-day supply</li> <li>Restasis Multidose®: 5.5 mL dispensed as a 25-day supply</li> <li>Tyrvaya™: 8.4 mL dispensed as a 30-day supply</li> <li>Verkazia®: 240 vials dispensed as a 30-day supply</li> </ul>
<b>Beta Blockers – Ophthalmic</b>		
betaxolol Betoptic S® carteolol Combigan® <sup>BLTG</sup> Istalol® levobunolol timolol maleate solution (gen Istalol®) timolol maleate gel (gen Timoptic-XE®)	Betimol® brimonidine/timolol (gen Combigan®) timolol maleate (gen Timoptic® and Timoptic® Ocudose®) Timoptic® Timoptic® Ocudose® Timoptic-XE®	

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XII. Ophthalmics</b>		
<b>Fluoroquinolones – Ophthalmic <sup>ST</sup></b>		
ciprofloxacin moxifloxacin (gen Vigamox®) ofloxacin	Besivance® Ciloxan® gatifloxacin levofloxacin moxifloxacin (gen Moxeza®) Ocuflox® Vigamox® Zymaxid®	<p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>For patients 21 years or younger, attempt treatment with a non-fluoroquinolone ophthalmic antibiotic before progressing to a fluoroquinolone ophthalmic product</li> <li>Examples of Non-Fluoroquinolone Ophthalmic Antibiotics <ul style="list-style-type: none"> <li>AK-Poly-Bac eye ointment</li> <li>bacitracin-polymyxin eye ointment</li> <li>erythromycin eye ointment</li> <li>Gentak® (3 mg/gm eye ointment, 3 mg/mL eye drops)</li> <li>gentamicin (3 mg/gm eye ointment, 3 mg/mL eye drops)</li> <li>neomycin-polymyxin-gramicidin eye drops</li> <li>polymyxin B-TMP eye drops</li> <li>Romycin® eye ointment</li> <li>sulfacetamide 10% eye drops</li> <li>Sulfamide® 10% eye drops</li> <li>tobramycin 0.3% eye drops</li> <li>Tobrasol™ 0.3% eye drops</li> </ul> </li> </ul>
<b>Non-Steroidal Anti-Inflammatory Drugs (NSAIDS) – Ophthalmic</b>		
diclofenac flurbiprofen Ilevro® ketorolac ketorolac LS	Acular® Acular LS® Acuvail® bromfenac BromSite® Nevanac® Prolensa®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XII. Ophthalmics</b>		
<b>Prostaglandin Agonists – Ophthalmic</b>		
latanoprost	bimatoprost Lumigan® Rocklatan™ tafluprost (gen Zioptan®) Travatan Z® travoprost (gen Travatan Z®) Xalatan® Xelpros™ Vyzulta™ Zioptan®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XIII. Otics</b>		
<b>Fluoroquinolones – Otic</b>		
Cipro HC® Ciprodex® <small>BLTG</small> ofloxacin	ciprofloxacin ciprofloxacin/dexamethasone (gen Ciprodex®) ciprofloxacin/fluocinolone (gen Otovel™) Otovel™	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XIV. Renal and Genitourinary</b>		
<b>Alpha Reductase Inhibitors for BPH</b>		
finasteride	Avodart® dutasteride dutasteride/tamsulosin Entadfi™ Jalyn® Proscar®	
<b>Antihyperuricemics</b>		
allopurinol 100 mg, 300 mg colchicine tablet febuxostat probencid probencid/colchicine	allopurinol 200 mg colchicine capsule Colcrys Gloperba® Mitigare® Uloric® Zyloprim®	
<b>Cystine Depleting Agents <small>CC</small></b>		
Cystagon®	Procysbi® <small>ST</small>	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul> <p><b>STEP THERAPY (ST)</b></p> <ul style="list-style-type: none"> <li>• Requires a trial with Cystagon immediate-release capsules</li> </ul>
<b>Phosphate Binders/Regulators</b>		
calcium acetate Renvela® tablet <small>BLTG</small> sevelamer HCl (gen Renagel)	Auryxia™ Fosrenol® lanthanum carbonate Phoslyra® Renvela® powder pack sevelamer carbonate powder, tablet (gen Renvela) Velphoro®	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XIV. Renal and Genitourinary</b>		
<b>Selective Alpha Adrenergic Blockers</b>		
alfuzosin tamsulosin	Flomax® Rapaflo® silodosin	
<b>Urinary Tract Antispasmodics</b>		
fesoterodine ER (gen Toviaz®) oxybutynin oxybutynin ER <small>DO, 1</small> solifenacin Toviaz® <small>DO</small>	darifenacin Detrol® Detrol LA® <small>DO</small> Ditropan XL® flavoxate Gelnique® Gemtesa® Myrbetriq® <small>DO</small> Myrbetriq® solution <small>F/Q/D</small> Oxytrol® tolterodine tolterodine ER trospium trospium ER Vesicare® <small>DO</small> Vesicare® LS	<p><b>DOSE OPTIMIZATION (DO)</b></p> <ul style="list-style-type: none"> <li>See Dose Optimization Chart for affected strengths</li> </ul> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <ul style="list-style-type: none"> <li>Myrbetriq® solution; limited to a 30-day supply</li> </ul>

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf) 56

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XV. Respiratory</b>		
<b>Anticholinergics / COPD Agents</b>		
Anoro Ellipta® Atrovent HFA® Bevespi® Aerosphere® Combivent Respimat® Incruse Ellipta® <sup>1</sup> ipratropium ipratropium / albuterol roflumilast (gen Daliresp®) <sup>1</sup> Spiriva® HandiHaler® Spiriva Respimat® Stiolto Respimat® Trelegy Ellipta® <sup>1</sup> Tudorza Pressair® <sup>1</sup>	Breztri™ Aerosphere Daliresp® Duaklir® Pressair Lonhala® Magnair® Yupelri®	
<b>Antihistamines – Intranasal</b>		
azelastine olopatadine	Patanase®	
<b>Antihistamines – Second Generation</b>		
cetirizine OTC tablet cetirizine OTC syrup/solution 1mg/ 1mL fexofenadine OTC tablet <sup>1</sup> levocetirizine tablet loratadine OTC	cetirizine OTC chewable cetirizine OTC syrup/solution 5 mg/5 mL cetirizine-D OTC Clarinex® Clarinex-D® desloratadine (gen Clarinex®) levocetirizine solution loratadine-D OTC	
<b>Beta2 Adrenergic Agents – Inhaled Long-Acting <sup>CC, F/Q/D</sup></b>		
arformoterol (gen Brovana®) <sup>1</sup> formoterol (gen Perforomist®) Serevent Diskus®	Brovana® Perforomist® Striverdi Respimat®	<b>CLINICAL CRITERIA (CC)</b> PA is required for all new long-acting beta agonist prescriptions for beneficiaries under FDA- or compendia-supported age as indicated:

1 = Preferred as of 08/03/2023

2 = Non-Preferred as of 08/03/2023

Standard PA fax form: [https://newyork.fhsc.com/downloads/providers/NYRx\\_PDP\\_PA\\_Fax\\_Standardized.pdf](https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Fax_Standardized.pdf)

57

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters																
<b>XV. Respiratory</b>																		
		<table border="1"> <tr> <td>Brovana® / arformoterol</td> <td>≥ 18 years</td> </tr> <tr> <td>Perforomist® / formoterol</td> <td>≥ 18 years</td> </tr> <tr> <td>Serevent Diskus®</td> <td>≥ 4 years</td> </tr> <tr> <td>Striverdi Respimat®</td> <td>≥ 18 years</td> </tr> </table> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <p>Maximum units per 30 days</p> <table border="1"> <tr> <td>Brovana® / arformoterol</td> <td>60 units (1 carton of 60 vials or 120 mL)</td> </tr> <tr> <td>Perforomist® / formoterol</td> <td>60 units (1 carton of 60 vials or 120 mL)</td> </tr> <tr> <td>Serevent Diskus®</td> <td>1 diskus (60 blisters)</td> </tr> <tr> <td>Striverdi Respimat®</td> <td>1 unit (one cartridge and one Respimat inhaler)</td> </tr> </table>	Brovana® / arformoterol	≥ 18 years	Perforomist® / formoterol	≥ 18 years	Serevent Diskus®	≥ 4 years	Striverdi Respimat®	≥ 18 years	Brovana® / arformoterol	60 units (1 carton of 60 vials or 120 mL)	Perforomist® / formoterol	60 units (1 carton of 60 vials or 120 mL)	Serevent Diskus®	1 diskus (60 blisters)	Striverdi Respimat®	1 unit (one cartridge and one Respimat inhaler)
Brovana® / arformoterol	≥ 18 years																	
Perforomist® / formoterol	≥ 18 years																	
Serevent Diskus®	≥ 4 years																	
Striverdi Respimat®	≥ 18 years																	
Brovana® / arformoterol	60 units (1 carton of 60 vials or 120 mL)																	
Perforomist® / formoterol	60 units (1 carton of 60 vials or 120 mL)																	
Serevent Diskus®	1 diskus (60 blisters)																	
Striverdi Respimat®	1 unit (one cartridge and one Respimat inhaler)																	
<b>Beta2 Adrenergic Agents – Inhaled Short-Acting</b>																		
albuterol nebulizer solution Ventolin HFA® <span style="color: red; text-decoration: underline;">BLTG</span>	albuterol HFA levalbuterol solution levalbuterol HFA ProAir® Digihaler™ ProAir® RespiClick Proventil HFA® Xopenex HFA®																	
<b>Corticosteroids – Inhaled F/Q/D</b>																		
Asmanex® Twisthaler Flovent Diskus® Flovent HFA® <span style="color: red; text-decoration: underline;">BLTG</span> Pulmicort® Flexhaler	Alvesco® ArmonAir® Digihaler® Arnuity Ellipta® Asmanex® HFA fluticasone HFA (gen Flovent® HFA) QVAR RediHaler®	<table border="1"> <tr> <td colspan="2"><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></td> </tr> <tr> <td>Alvesco® 80 mcg</td> <td>1 inhaler every 30 days</td> </tr> <tr> <td>Alvesco® 160 mcg</td> <td>1 inhaler every 30 days. Up to 1 inhaler every 15 days with previous oral corticosteroid use.</td> </tr> <tr> <td>ArmonAir® Digihaler®</td> <td>1 inhaler every 30 days</td> </tr> <tr> <td>Arnuity Ellipta</td> <td>1 inhaler every 30 days</td> </tr> <tr> <td>Asmanex® 110 mcg</td> <td>1 inhaler every 30 days</td> </tr> <tr> <td>Asmanex® 220 mcg (30 units)</td> <td>1 inhaler every 30 days</td> </tr> </table>	<b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b>		Alvesco® 80 mcg	1 inhaler every 30 days	Alvesco® 160 mcg	1 inhaler every 30 days. Up to 1 inhaler every 15 days with previous oral corticosteroid use.	ArmonAir® Digihaler®	1 inhaler every 30 days	Arnuity Ellipta	1 inhaler every 30 days	Asmanex® 110 mcg	1 inhaler every 30 days	Asmanex® 220 mcg (30 units)	1 inhaler every 30 days		
<b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b>																		
Alvesco® 80 mcg	1 inhaler every 30 days																	
Alvesco® 160 mcg	1 inhaler every 30 days. Up to 1 inhaler every 15 days with previous oral corticosteroid use.																	
ArmonAir® Digihaler®	1 inhaler every 30 days																	
Arnuity Ellipta	1 inhaler every 30 days																	
Asmanex® 110 mcg	1 inhaler every 30 days																	
Asmanex® 220 mcg (30 units)	1 inhaler every 30 days																	

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters	
<b>XV. Respiratory</b>			
		Asmanex® 220 mcg (60 units)	1 inhaler every 30 days. Up to 1 inhaler every 15 days with previous oral corticosteroid use.
		Asmanex® 220 mcg (120 units)	1 inhaler every 60 days. Up to 1 inhaler every 30 days with previous oral corticosteroid use.
		Asmanex® HFA 100 mcg	1 inhaler every 30 days
		Asmanex® HFA 200 mcg	1 inhaler every 30 days
		Flovent Diskus® 50 mcg, 100 mcg	1 diskus every 30 days
		Flovent Diskus® 250 mcg	1 diskus every 15 days. Up to 1 diskus every 7 days with previous oral corticosteroid use.
		Flovent HFA® 44 mcg, 110 mcg	1 inhaler every 30 days
		Flovent HFA® 220 mcg	1 inhaler every 30 days. Up to 1 inhaler every 15 days with previous oral corticosteroid use.
		Pulmicort 90 mcg	1 inhaler every 30 days
		Pulmicort 180 mcg	1 inhaler every 15 days
		QVAR® RediHaler™ 40 mcg	1 inhaler every 30 days
		QVAR® RediHaler™ 80 mcg	1 inhaler every 15 days

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters																											
<b>XV. Respiratory</b>																													
<b>Corticosteroid/Beta2 Adrenergic Agent (Long-Acting) Combinations – Inhaled</b> <span style="color: red;">CC, F/Q/D</span>																													
Advair Diskus® <span style="color: red;">BLTG</span> Dulera® Symbicort® <span style="color: red;">BLTG</span>	Advair HFA® AirDuo® Digihaler® AirDuo™ RespiClick® Breo Ellipta® budesonide/formoterol (gen Symbicort) fluticasone-salmeterol (gen AirDuo™ RespiClick®) fluticasone-salmeterol (gen Advair Diskus®) fluticasone-salmeterol (gen Advair HFA™) fluticasone-vilanterol (gen Breo Ellipta®)	<p><b>CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>PA is required for all new long-acting beta agonist prescriptions for beneficiaries under FDA-or compendia-supported age as indicated:</li> </ul> <table border="1"> <tr><td>Advair Diskus®</td><td>≥ 4 years</td></tr> <tr><td>Advair HFA®</td><td>≥ 12 years</td></tr> <tr><td>AirDuo™ RespiClick® &amp; Digihaler®</td><td>&gt; 12 years</td></tr> <tr><td>Dulera® 100 mcg and 200 mcg</td><td>≥ 12 years</td></tr> <tr><td>Dulera® 50 mcg</td><td>≥ 4 years</td></tr> <tr><td>fluticasone-salmeterol</td><td>≥ 4 years</td></tr> <tr><td>budesonide-formoterol (Symbicort®) 80/4.5 mcg</td><td>≥ 4 years</td></tr> <tr><td>budesonide-formoterol (Symbicort®) 160/4.5 mcg</td><td>≥ 12 years</td></tr> <tr><td>fluticasone/vilanterol (Breo Ellipta®)</td><td>≥ 18 years</td></tr> </table> <p><b>FREQUENCY/QUANTITY/DURATION (F/Q/D)</b></p> <table border="1"> <tr><td>Advair Diskus®</td><td rowspan="5" style="vertical-align: middle; text-align: center;">One inhaler/diskus every 30 days</td></tr> <tr><td>Advair HFA®</td></tr> <tr><td>AirDuo™ RespiClick® &amp; Digihaler®</td></tr> <tr><td>fluticasone-salmeterol</td></tr> <tr><td>fluticasone/vilanterol (Breo Ellipta®)</td></tr> <tr><td>Budesonide/formoterol (Symbicort®)</td><td rowspan="2" style="vertical-align: middle; text-align: center;">Up to 8 inhalers every 180 days</td></tr> <tr><td>Dulera®</td></tr> </table>	Advair Diskus®	≥ 4 years	Advair HFA®	≥ 12 years	AirDuo™ RespiClick® & Digihaler®	> 12 years	Dulera® 100 mcg and 200 mcg	≥ 12 years	Dulera® 50 mcg	≥ 4 years	fluticasone-salmeterol	≥ 4 years	budesonide-formoterol (Symbicort®) 80/4.5 mcg	≥ 4 years	budesonide-formoterol (Symbicort®) 160/4.5 mcg	≥ 12 years	fluticasone/vilanterol (Breo Ellipta®)	≥ 18 years	Advair Diskus®	One inhaler/diskus every 30 days	Advair HFA®	AirDuo™ RespiClick® & Digihaler®	fluticasone-salmeterol	fluticasone/vilanterol (Breo Ellipta®)	Budesonide/formoterol (Symbicort®)	Up to 8 inhalers every 180 days	Dulera®
Advair Diskus®	≥ 4 years																												
Advair HFA®	≥ 12 years																												
AirDuo™ RespiClick® & Digihaler®	> 12 years																												
Dulera® 100 mcg and 200 mcg	≥ 12 years																												
Dulera® 50 mcg	≥ 4 years																												
fluticasone-salmeterol	≥ 4 years																												
budesonide-formoterol (Symbicort®) 80/4.5 mcg	≥ 4 years																												
budesonide-formoterol (Symbicort®) 160/4.5 mcg	≥ 12 years																												
fluticasone/vilanterol (Breo Ellipta®)	≥ 18 years																												
Advair Diskus®	One inhaler/diskus every 30 days																												
Advair HFA®																													
AirDuo™ RespiClick® & Digihaler®																													
fluticasone-salmeterol																													
fluticasone/vilanterol (Breo Ellipta®)																													
Budesonide/formoterol (Symbicort®)	Up to 8 inhalers every 180 days																												
Dulera®																													

## NYRx, the Medicaid Pharmacy Program Preferred Drug List

Preferred Drugs	Non-Preferred Drugs	Prior Authorization/Coverage Parameters
<b>XV. Respiratory</b>		
<b>Corticosteroids – Intranasal</b>		
fluticasone	azelastine-fluticasone (gen Dymista®) Beconase AQ® <small>CC</small> Dymista® flunisolide mometasone Omnaris® QNASL® <small>CC</small> Ryaltris® Xhance™ Zetonna®	<b>CLINICAL CRITERIA (CC)</b> <ul style="list-style-type: none"> <li>• Clinical consideration in regard to drug interactions will be given to patients with HIV/AIDS diagnosis or antiretroviral therapy in history</li> </ul>
<b>Leukotriene Modifiers</b>		
montelukast tablet, chew tab <small>ST</small>	Accolate® montelukast granules Singulair® <small>ST</small> zafirlukast	<b>STEP THERAPY (ST)</b> <ul style="list-style-type: none"> <li>• For non-asthmatic patients, trial of intranasal corticosteroid or a 2nd generation oral antihistamine before montelukast (Singulair®)</li> </ul>

# NYRx, the Medicaid Pharmacy Program Preferred Drug List

## NYS Medicaid NYRx Clinical Drug Review Program (CDRP)

The Clinical Drug Review Program (CDRP) is aimed at ensuring specific drugs are utilized in a medically appropriate manner.

Under the CDRP, certain drugs require prior authorization because there may be specific safety issues, public health concerns, the potential for fraud and abuse or the potential for significant overuse and misuse.

### Prior Authorization

Prior authorization for some drugs subject to the CDRP must be obtained through a representative at the clinical call center. For some drugs subject to the CDRP, only prescribers, not their authorized agents, can initiate the prior authorization process.

Please be prepared to respond to a series of questions that identify prescriber, patient, and reason for prescribing drug, and to fax clinical documentation upon request. Clinical guidelines for the CDRP as well as prior authorization worksheets are available online at [https://newyork.fhsc.com/providers/CDRP\\_about.asp](https://newyork.fhsc.com/providers/CDRP_about.asp).

The following drugs are subject to the Clinical Drug Review Program:

- fentanyl mucosal agents: [https://newyork.fhsc.com/providers/CDRP\\_fentanyl\\_mucosal\\_agents.asp](https://newyork.fhsc.com/providers/CDRP_fentanyl_mucosal_agents.asp)
- palivizumab (Synagis®): [https://newyork.fhsc.com/providers/CDRP\\_synagis.asp](https://newyork.fhsc.com/providers/CDRP_synagis.asp)
- sodium oxybate products (Xyrem®, Xywav™): [https://newyork.fhsc.com/providers/CDRP\\_xyrem.asp](https://newyork.fhsc.com/providers/CDRP_xyrem.asp)
- somatropin (Serostim®): [https://newyork.fhsc.com/providers/CDRP\\_serostim.asp](https://newyork.fhsc.com/providers/CDRP_serostim.asp)

The following drug classes are subject to the Clinical Drug Review Program and are also included on the Preferred Drug List:

- Anabolic Steroids: [https://newyork.fhsc.com/providers/CDRP\\_anabolic\\_steroids.asp](https://newyork.fhsc.com/providers/CDRP_anabolic_steroids.asp)
- Growth Hormones for 18 years and older: [https://newyork.fhsc.com/providers/CDRP\\_growth\\_hormones.asp](https://newyork.fhsc.com/providers/CDRP_growth_hormones.asp)

## NYS Medicaid NYRx Drug Utilization Review (DUR) Program

Frequency/Quantity/Duration (F/Q/D) Program and Step Therapy parameters are implemented to ensure clinically appropriate and cost effective use of these drugs and drug classes.

For additional Step Therapy and Frequency/Quantity/Duration parameters for drugs and drug classes that are also included on the Preferred Drug List (PDL), please see pages 3 through 60.

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Corticotropin (Acthar®, ACTH injectable, Cortrophin®)	<p>Requires trial of first-line therapy for all FDA-approved indications, other than infantile spasms.</p> <p><b>Note:</b> It is first line therapy for infantile spasms in children less than 2 years of age – step therapy not required.</p>	<p><b>QUANTITY LIMITS:</b></p> <ul style="list-style-type: none"> <li>• Infantile spasms – 30 mL (six 5 mL vials)</li> <li>• Multiple sclerosis – 35 mL (seven 5 mL vials)</li> </ul> <p><b>DURATION LIMITS:</b></p> <ul style="list-style-type: none"> <li>• Infantile spasms – 4 weeks; indicated for &lt; 2 years of age</li> <li>• Multiple sclerosis – 5 weeks</li> <li>• Rheumatic disorders – 5 weeks</li> <li>• Dermatologic conditions – 5 weeks</li> <li>• Allergic states (serum sickness) – 5 weeks</li> </ul>	<ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved or compendia-supported indication</li> <li>• Not covered for diagnostic purposes</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Corticotropin (Acthar®, ACTH injectable, Cortrophin®) (continued)		FDA Indication	First Line Therapy

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Anabolic Steroids – Injectable <ul style="list-style-type: none"> <li>• Depo-Testosterone®</li> <li>• testosterone cypionate*</li> <li>• testosterone enanthate</li> <li>• Xyosted®</li> </ul>		<ul style="list-style-type: none"> <li>• Limitations for anabolic steroid products is based on approved FDA labeled daily dosing and documented diagnosis not to exceed a 90-day supply (30-day supply for oxandrolone):</li> </ul>	*for additional parameters, see Cross-Sex Hormones section below.
Anabolic Steroids – Oral <ul style="list-style-type: none"> <li>• Jatenzo®</li> <li>• Methitest®</li> <li>• Oxandrolone</li> <li>• Tlando®</li> </ul>		<ul style="list-style-type: none"> <li>• Xyosted® is limited to no more than 3 boxes for 90 days (1 box per 30 days)</li> <li>• Initial duration limit of 3 months (for all products except oxandrolone), requiring documented follow-up monitoring for response and/or adverse effects before continuing treatment</li> <li>• Duration limit of 6 months for delayed puberty</li> <li>• Duration limit of 1 month for all uses of oxandrolone products</li> </ul>	
Anti-Diabetic agents (not on the PDL) <ul style="list-style-type: none"> <li>• acarbose (Precose®)</li> <li>• glimepiride</li> <li>• glipizide (Glucotrol XL®)</li> <li>• glyburide</li> <li>• glyburide, micronized</li> <li>• miglitol</li> <li>• nateglinide</li> <li>• pramlintide (Symlin®)</li> <li>• repaglinide</li> <li>• repaglinide/metformin</li> </ul>	<ul style="list-style-type: none"> <li>• Requires a trial with metformin with or without insulin prior to initiating other antidiabetic agents unless there is a documented contraindication.</li> <li>• Clinical editing to allow patients with a diagnosis of gestational diabetes to receive glyburide without a trial of metformin first.</li> </ul>		

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Anti-Diarrheal Agents <ul style="list-style-type: none"> <li>• alosetron (Lotronex®)</li> <li>• crofelemer (Mytesi®)</li> <li>• eluxadoline (Viberzi®)</li> <li>• telotristat (Xermelo®)</li> </ul>	<ul style="list-style-type: none"> <li>• Irritable Bowel Syndrome w/Diarrhea               <ul style="list-style-type: none"> <li>– Trial of eluxadoline and rifaximin prior to alosetron.</li> </ul> </li> <li>• Symptomatic relief of non-infectious diarrhea in patients with HIV/AIDS on anti-retroviral therapy               <ul style="list-style-type: none"> <li>– Trial with an alternative anti-diarrheal agent.</li> </ul> </li> <li>• Carcinoid Syndrome               <ul style="list-style-type: none"> <li>– Trial with and concurrent use with a somatostatin analog</li> </ul> </li> </ul>		<ul style="list-style-type: none"> <li>• Confirmation of FDA-approved or compendia-supported indication.</li> </ul>
Anti-Fungals, Topical – for Onychomycosis <ul style="list-style-type: none"> <li>• ciclopirox 8% solution</li> <li>• Jublia®</li> <li>• tavaborole (Kerydin®)</li> </ul>	<ul style="list-style-type: none"> <li>• Trial with an oral antifungal agent* prior to use of ciclopirox 8% solution                *terbinafine (Lamisil®) tablets; griseofulvin (Gris PEG®) oral suspension, ultramicrogranulated tablets; micronized tablets; itraconazole (Sporanox®,) tablets, oral solution               <ul style="list-style-type: none"> <li>• Trial with ciclopirox 8% solution prior to the use of other topical antifungals [efinaconazole (Jublia®) or tavaborole (Kerydin®)]</li> </ul> </li> </ul>		
Anti-Malarias <ul style="list-style-type: none"> <li>• chloroquine</li> <li>• hydroxychloroquine</li> </ul>			<ul style="list-style-type: none"> <li>• Confirm FDA-approved or compendia-supported use</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Anti-Retroviral (ARV) Interventions		<b>QUANTITY LIMITS:</b> <ul style="list-style-type: none"> <li>Limit ARV active ingredient duplication</li> <li>Limit ARV utilization to a maximum of five products concurrently - excluding boosting with ritonavir (dose limit 600 mg or less) or cobicistat</li> <li>Limit Protease Inhibitor utilization to a maximum of two products concurrently</li> <li>Limit Integrase inhibitor utilization to a maximum of one product concurrently</li> </ul>	<ul style="list-style-type: none"> <li>Require confirmation of FDA-approved or compendia-supported use</li> <li>Point-of-service edit for antiretroviral / non-antiretroviral combinations to be avoided: <a href="https://newyork.fhsc.com/downloads/providers/NYRx_PDP_reference_Antiretroviral_NonAntiretroviral_Drug2Drug_Interactions.pdf">https://newyork.fhsc.com/downloads/providers/NYRx_PDP_reference_Antiretroviral_NonAntiretroviral_Drug2Drug_Interactions.pdf</a></li> <li>Point-of-service edit for antiretroviral / antiretroviral combinations to be avoided: <a href="https://newyork.fhsc.com/downloads/providers/NYRx_PDP_reference_Antiretroviral_Antiretroviral_Drug2Drug_Interactions.pdf">https://newyork.fhsc.com/downloads/providers/NYRx_PDP_reference_Antiretroviral_Antiretroviral_Drug2Drug_Interactions.pdf</a></li> </ul>
Benlysta® (belimumab)	<ul style="list-style-type: none"> <li>Trial of a disease-modifying anti-rheumatic drug (DMARD) prior to treatment with an immunomodulator</li> </ul>		<ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul>
biotin			<ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul>
Atopic Dermatitis Agents • crisaborole (Eucrisa®) • ruxolitinib (Opzelura™)	<ul style="list-style-type: none"> <li>Trial with a medium or high potency prescription topical steroid within the last 3 months</li> </ul>	<b>QUANTITY LIMITS:</b> <ul style="list-style-type: none"> <li>100 gm/30 days (crisaborole)</li> <li>240 gm/30 days (ruxolitinib)</li> </ul>	<ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> <li>ruxolitinib: age 12 years +</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Benzodiazepine agents – oral <ul style="list-style-type: none"> <li>• alprazolam (Xanax®, Xanax® XR)</li> <li>• chlordiazepoxide</li> <li>• chlordiazepoxide/amitriptyline</li> <li>• clonazepam (Klonopin®)</li> <li>• clorazepate</li> <li>• diazepam (Valium®)</li> <li>• lorazepam (Ativan®, Lorazepam Intensol®, Loreev XR™)</li> <li>• oxazepam</li> </ul>	Generalized Anxiety Disorder (GAD) or Social Anxiety Disorder (SAD) <ul style="list-style-type: none"> <li>• Require trial with a Selective-Serotonin Reuptake Inhibitor (SSRI) or a Serotonin-Norepinephrine Reuptake Inhibitor (SNRI) prior to initial benzodiazepine prescription</li> <li>• Panic Disorder requires concurrent therapy with an antidepressant (SSRI, SNRI, or Tricyclic antidepressant [TCA]).</li> </ul> Skeletal muscle spasms <ul style="list-style-type: none"> <li>• Require trial with a skeletal muscle relaxant prior to a benzodiazepine</li> </ul>	<b>DURATION LIMIT:</b> <ul style="list-style-type: none"> <li>• For Insomnia: 30 consecutive days</li> <li>• For Panic Disorder: 30 consecutive days</li> </ul>	<ul style="list-style-type: none"> <li>• Require confirmation of FDA-approved or compendia-supported use</li> <li>• PA required for initiation of benzodiazepine therapy in patients currently on opioid or oral buprenorphine therapy</li> <li>• PA required for any additional oral benzodiazepine prescription in patients currently on benzodiazepine therapy</li> <li>• PA required when greater than a 14-day supply of a benzodiazepine is prescribed for someone on a CNS stimulant</li> </ul>
Constipation Agents <ul style="list-style-type: none"> <li>• linaclootide (Linzess®)</li> <li>• lubiprostone (Amitiza®)</li> <li>• methylnaltrexone (Relistor®)</li> <li>• naldemedine (Symproic®)</li> <li>• naloxegol (Movantik®)</li> <li>• plecanatide (Trulance®)</li> <li>• prucalopride (Motegrity™)</li> <li>• tenapanor (Ibsrela®)</li> </ul>	Opioid Induced Constipation (OIC) and Chronic Idiopathic Constipation (CIC) <ul style="list-style-type: none"> <li>• Trial with an osmotic laxative, a stimulant laxative, and a stool softener prior to use.</li> </ul> Irritable Bowel Syndrome w/ Constipation (IBS-C) <ul style="list-style-type: none"> <li>• Trial with a bulking agent and an osmotic laxative within 89 days of use.</li> </ul>	<b>QUANTITY LIMIT:</b> <ul style="list-style-type: none"> <li>• linaclootide, naldemedine, naloxegol, plecanatide: 1 tablet/day</li> <li>• lubiprostone: 2 capsules/day</li> <li>• methylnaltrexone: 1 vial or syringe/day, 4 kits/28 days</li> <li>• prucalopride: 2 mg/day max; 1 tablet per day</li> <li>• tenapanor 2 tablets/day</li> </ul>	<ul style="list-style-type: none"> <li>• Confirmation of FDA-approved or compendia-supported indication.</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Cross-Sex Hormones <ul style="list-style-type: none"> <li>• conjugated estrogens</li> <li>• estradiol</li> <li>• testosterone cypionate</li> <li>• testosterone enanthate (Xyoste<sup>TM</sup>)</li> <li>• testosterone gel 1.62% (AndroGel<sup>®</sup>)*</li> <li>• testosterone patch*</li> </ul>			<ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved or compendia-supported indication</li> <li>• For diagnosis of gender dysphoria please refer to July 2020 edition of the Medicaid Update: <a href="https://www.health.ny.gov/health_care/medicaid/program/update/2020/no12_2020-07.htm#transgender">https://www.health.ny.gov/health_care/medicaid/program/update/2020/no12_2020-07.htm#transgender</a></li> </ul>
*Subject to Anabolic Steroids – Topical PDL class criteria			
dextromethorphan / quinidine (Nuedexta <sup>®</sup> )		<b>QUANTITY LIMIT:</b> <ul style="list-style-type: none"> <li>• 2 capsules per day; 60 units per 30 days</li> </ul> <b>DURATION LIMIT:</b> <ul style="list-style-type: none"> <li>• 90 days of therapy</li> </ul>	For patients ≥ 18 years of age: <ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul>
Diabetic Test Strips		<b>QUANTITY LIMIT:</b> <ul style="list-style-type: none"> <li>• Type I DM – max 300 test strips per 30-day supply</li> <li>• Type II DM – max 100 test strips per 30-day supply</li> </ul>	<ul style="list-style-type: none"> <li>• Preferred diabetic supply program <a href="https://newyork.fhsc.com/providers/diabeticsupplies.asp">https://newyork.fhsc.com/providers/diabeticsupplies.asp</a></li> </ul>
Direct Renin Inhibitors <ul style="list-style-type: none"> <li>• aliskiren</li> <li>• Tekturna<sup>®</sup></li> <li>• Tekturna<sup>®</sup> HCT</li> </ul>	Trial of product containing either an ACE inhibitor or an ARB prior to initiating preferred DRI		

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
dronabinol (Marinol®)	<p>Step therapy for beneficiaries with HIV/AIDS, or cancer, AND eating disorder:</p> <ul style="list-style-type: none"> <li>• Trial with megestrol acetate suspension prior to dronabinol</li> </ul> <p>Step therapy for beneficiaries with diagnosis of cancer and nausea/vomiting:</p> <ul style="list-style-type: none"> <li>• Trial with a NYS Medicaid-preferred 5-HT3 receptor antagonist prior to dronabinol</li> </ul>		<ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul>
risdiplam (Evrysdi®)			<ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved indication</li> <li>• Confirm absence of advanced disease</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Fentanyl Transmucosal Agents <ul style="list-style-type: none"> <li>• Actiq® (lozenge)</li> <li>• Fentora® (buccal tablet)</li> </ul>		<b>QUANTITY LIMIT:</b> Actiq®, Fentora: <ul style="list-style-type: none"> <li>• 4 units per day, 120 units per 30 days</li> </ul> <b>DURATION LIMIT:</b> <ul style="list-style-type: none"> <li>• 90 days</li> <li>• Exemption for diagnosis of cancer, sickle cell disease, or hospice care</li> </ul>	<ul style="list-style-type: none"> <li>• Limited to a total of 4 opioid prescriptions every 30 days;</li> <li>• For opioid-naïve patients: limited to a 7 days' supply for all initial opioid prescriptions,</li> <li>• PA required for use if &gt; 90 MME (MME = morphine milligram equivalents) of opioid per day for management of non-acute pain (pain lasting &gt; 7 days).</li> <li>• PA required for initiation of opioid therapy for patients on established opioid dependence therapy</li> <li>• PA is required for initiation of opioid therapy in patients currently on benzodiazepine therapy</li> <li>• Exemption for diagnosis of cancer, sickle cell, or hospice care</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
HIV PrEP (Pre-Exposure Prophylaxis Agents): <ul style="list-style-type: none"> <li>• cabotegravir (Apretude)</li> <li>• emtricitabine/tenofovir disoproxil fumarate (Truvada®)</li> <li>• emtricitabine/tenofovir alafenamide (Descovy®)</li> </ul>			<ul style="list-style-type: none"> <li>• Prescribers or authorized agents are required to respond to a series of questions that identify the prescriber, the patient, and the reason for prescribing an HIV-1 PrEP agent.</li> <li>• Prescribers or authorized agents must indicate whether the HIV-1 PrEP agent has been prescribed for HIV pre-exposure prophylaxis (PrEP) or treatment of HIV/AIDS. If the agent has been prescribed for prophylaxis, the date of last negative HIV test must also be provided.</li> </ul>
ivermectin (oral)			<ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul>
Lidocaine patches <ul style="list-style-type: none"> <li>• Lidoderm®</li> <li>• ZTLido™</li> </ul>			<ul style="list-style-type: none"> <li>• Prescribers, or their authorized agents, are required to respond to a series of questions that identify the prescriber, the patient, and the reason for prescribing this drug.</li> <li>• Prescriptions can be written for a 30-day supply with up to 2 refills</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
<p>Lipid Lowering Agents:</p> <ul style="list-style-type: none"> <li>• alirocumab (Praluent®)</li> <li>• evolocumab (Repatha®)</li> <li>• lomitapide (Juxtapid®)</li> <li>• bempedoic acid (Nexletol™)</li> <li>• bempedoic acid/ezetimibe (Nexlizet™)</li> </ul>	<ul style="list-style-type: none"> <li>• Require trial of an HMG-CoA Reductase Inhibitors (statin) at maximum tolerated dosage</li> </ul>		<ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul> <p><b>PCSK-9 Inhibitors</b> (alirocumab [Praluent®], evolocumab [Repatha®]) and <b>ACL inhibitors</b> (Bempedoic acid [Nexletol], Bempedoic acid/ezetimibe [Nexlizet]):</p> <ul style="list-style-type: none"> <li>• Require concurrent statin therapy</li> </ul>
Methadone	<ul style="list-style-type: none"> <li>• Requires a trial of a long-acting opioid prior to initiation for the management of chronic non-cancer pain</li> </ul>	<p><b>QUANTITY LIMIT:</b></p> <ul style="list-style-type: none"> <li>• 12 units per day, 360 units per 30 days</li> <li>• Exemption for diagnosis of cancer, hospice care, or sickle cell disease</li> </ul>	<ul style="list-style-type: none"> <li>• Confirm diagnosis of chronic non-cancer pain</li> <li>• Limited to a total of 4 opioid prescriptions every 30 days;</li> <li>• PA required for initiation of methadone for patients on established opioid dependence therapy</li> <li>• PA required for methadone prescriptions for patients currently on long-acting opioid therapy.</li> <li>• PA required for initiation of long-acting opioid therapy in opioid-naïve patients.</li> <li>• PA required for use if &gt; 90 MME (MME = morphine milligram equivalents) of opioid per day for management of non-acute pain (pain lasting &gt; 7 days). PA required for initiation of methadone therapy in patients currently on benzodiazepine therapy</li> <li>• Exemption for diagnosis of cancer, sickle cell, or hospice care</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Metoclopramide nasal spray (Gimoti™)			<ul style="list-style-type: none"> <li>Metoclopramide nasal spray confirm diagnosis of diabetes</li> </ul>
olanzapine / fluoxetine (Symbyax®)	<ul style="list-style-type: none"> <li>When prescribing for the treatment of major depressive disorder (MDD) in the absence of other psychiatric comorbidities, trial with at least one different antidepressant agent is required</li> </ul>		<ul style="list-style-type: none"> <li>PA is required for the initial prescription for beneficiaries younger than 10 years</li> </ul>
Ovulation Enhancing Drugs <ul style="list-style-type: none"> <li>bromocriptine</li> <li>clomiphene</li> <li>letrozole</li> <li>tamoxifen</li> </ul>			<ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication and Medicaid covered indication</li> <li>Refer to <a href="https://www.health.ny.gov/health_care/medicaid/program/update/2019/2019-06.htm#ovulation">https://www.health.ny.gov/health_care/medicaid/program/update/2019/2019-06.htm#ovulation</a></li> </ul>
Oxazolidinone Antibiotics <ul style="list-style-type: none"> <li>linezolid (Zyvox®)</li> <li>tedizolid (Sivextro®)</li> </ul>			<ul style="list-style-type: none"> <li>Please be prepared to respond to a series of questions that identify the prescriber, the patient, and the reason for prescribing this drug.</li> <li>Please be prepared to fax clinical documentation upon request.</li> </ul>
Pubertal Suppressants <ul style="list-style-type: none"> <li>goserelin acetate</li> <li>leuprolide acetate</li> <li>nafarelin acetate</li> </ul>			<ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> <li>Refer to <a href="https://www.health.ny.gov/health_care/medicaid/program/update/2017/2017-01.htm#transgender">https://www.health.ny.gov/health_care/medicaid/program/update/2017/2017-01.htm#transgender</a> for Transgender Related Care and Services Update</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Rosacea Agents <ul style="list-style-type: none"> <li>• azelaic acid (Finacea®)</li> <li>• brimonidine gel pump</li> <li>• ivermectin</li> <li>• oxymetazoline HCl (Rhofade®)</li> <li>• doxycycline</li> </ul>	<ul style="list-style-type: none"> <li>• Trial with topical metronidazole product.</li> </ul>		<ul style="list-style-type: none"> <li>• Confirmation of FDA-approved or compendia-supported indication</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Spravato® (esketamine)	<ul style="list-style-type: none"> <li>Treatment Resistant Depression: trial of at least two oral antidepressants</li> </ul>		<ul style="list-style-type: none"> <li>Confirm diagnosis of FDA approved indication for patients ≥18 years of age</li> <li>Confirm concurrent use of an FDA approved antidepressant</li> <li>Before initiating esketamine nasal spray (Spravato), prescribers must attest that they have obtained a baseline score using a validated clinical assessment tool for depression (e.g., HAMD17, QIDS-C16C, MADRS).</li> <li>After the initiation of esketamine nasal spray (Spravato) therapy, every six months prescribers must attest that esketamine nasal spray (Spravato) has resulted in an improvement of depressive symptoms (from baseline) using the same baseline clinical assessment tool for depression (e.g., HAMD17, QIDS-C16C, MADRS). The esketamine worksheet can be accessed at: <a href="https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Worksheet_Prescribers_Spravato.docx">https://newyork.fhsc.com/downloads/providers/NYRx_PDP_PA_Worksheet_Prescribers_Spravato.docx</a></li> </ul>
tasimelteon (Hetlioz®, Hetlioz® LQ)		<b>QUANTITY LIMIT:</b> <ul style="list-style-type: none"> <li>One unit per day; 30 units per 30 days</li> </ul>	<ul style="list-style-type: none"> <li>Confirm diagnosis of FDA-approved or compendia-supported indication</li> </ul>

Drug / Class Name	Step Therapy (ST) Parameters	Frequency / Quantity / Duration (F/Q/D) Parameters	Additional / Alternate Parameter(s)
Parathyroid Hormone Analogs <ul style="list-style-type: none"> <li>• teriparatide (Forteo®)</li> <li>• Tymlos®</li> </ul>	<ul style="list-style-type: none"> <li>• Requires a trial with a preferred oral bisphosphonate</li> </ul>	<b>QUANTITY LIMIT:</b> <ul style="list-style-type: none"> <li>• One unit per 30-day period</li> </ul> <b>LIFETIME QUANTITY LIMIT:</b> <ul style="list-style-type: none"> <li>• 25 months' cumulative use of a PTH analog</li> </ul>	
Topical Compounded Prescriptions			<ul style="list-style-type: none"> <li>• Confirm diagnosis of FDA-approved or compendia-supported indication</li> <li>• For non-opioid pain management alternatives please visit: <a href="https://health.ny.gov/health_care/medicaid/program/opioid_management/docs/non_opioid_alternatives_to_pain_management.pdf">https://health.ny.gov/health_care/medicaid/program/opioid_management/docs/non_opioid_alternatives_to_pain_management.pdf</a></li> </ul>
Uterine Disorder Agents <ul style="list-style-type: none"> <li>• Oriahnn®</li> <li>• Myfembree®</li> </ul>		<b>QUANTITY LIMIT:</b> <ul style="list-style-type: none"> <li>• 28 days per 30-day period</li> </ul> <b>LIFETIME QUANTITY LIMIT:</b> <ul style="list-style-type: none"> <li>• 24 months cumulative use</li> </ul>	

For more information on DUR Program, please refer to [https://www.health.ny.gov/health\\_care/medicaid/program/dur/index.htm](https://www.health.ny.gov/health_care/medicaid/program/dur/index.htm).

## Medication Assisted Treatment (MAT) Formulary

Prior authorization will not be required for medications used for the treatment of substance use disorder prescribed according to generally accepted national professional guidelines for the treatment of a substance use disorder.

Effective 05/05/2022

### Medication Assisted Treatment (MAT) Formulary

\*\*Prior authorization will not be required when prescribed according to generally accepted national professional guidelines for the treatment of a substance use disorder. \*\*

Drugs	Coverage Parameters
<b>Opioid Antagonists</b>	
naloxone (syringe, vial) naltrexone Narcan® (nasal spray) naloxone nasal spray Kloxxado™ Zimhi™*	n/a
<b>Opioid Dependence Agents – Injectable</b>	
Vivitrol® Sublocade™	n/a
<b>Opioid Dependence Agents – Oral/Transmucosal F/Q/D</b>	
buprenorphine (tablet) buprenorphine/naloxone (tablet) Suboxone® (film) buprenorphine/naloxone (film) Zubsolv®	<p><b>QUANTITY LIMIT:</b></p> <ul style="list-style-type: none"> <li>• <b>buprenorphine sublingual (SL):</b> Six tablets dispensed as a 2-day supply; not to exceed 24 mg per day</li> <li>• <b>buprenorphine/ naloxone tablet and film (Suboxone®, Zubsolv®)</b> up to 5.7 mg/1.4 mg strength); Three sublingual tablets or films per day; maximum of 90 tablets or films dispensed as a 30-day supply, not to exceed 24 mg-6 mg of Suboxone, or its equivalent per day</li> <li>• <b>buprenorphine/naloxone tablet (Zubsolv® 8.6 mg/2.1 mg strength):</b> Maximum of 60 tablets dispensed as a 30-day supply</li> <li>• <b>buprenorphine/naloxone tablet (Zubsolv® 11.4 mg/2.9 mg strength):</b> Maximum of 30 tablets dispensed as a 30-day supply</li> </ul> <p><b>RELATED CLINICAL CRITERIA (CC)</b></p> <ul style="list-style-type: none"> <li>• PA required for initiation of <b>opioid therapy</b> for patients on established opioid dependence therapy</li> <li>• PA required for initiation of a <b>CNS stimulant</b> for patients established on opioid dependence therapy **</li> </ul>

## NYS Medicaid NYRx Brand Less Than Generic (BLTG) Program

On April 26, 2010, NYRx, the Medicaid Pharmacy Program, implemented a new cost-containment initiative, which promotes the use of certain multi-source brand name drugs when the cost of the brand name drug is less expensive than the generic equivalent.

In conformance with State Education Law, which intends that patients receive the lower cost alternative, brand name drugs included in this program:

- Do not require “Dispense as Written” (DAW) or “Brand Medically Necessary” on the prescription
- Have a generic copayment
- Are paid at the Brand Name Drug reimbursement rate or usual and customary price, whichever is lower (SMAC/FUL are not applied)
- Do not require a new prescription if the drug is removed from this program

### **Effective July 13, 2023:**

- Dexilant®, Oseni®, Protonix® suspension, Pylera®, Vascepa®, Viibryd® will be **added** to the program
- No products will be **removed** from the program

List of Brand Name Drugs included in this program**		
Advair Diskus®	EpiPen, Jr	<b>Pylera®</b>
Alphagan P® 0.15%	Firvanq®	Rapamune® solution
Amitiza®	Flovent® HFA	Renvela® tablet
Apriso®	Glumetza®	Restasis®
Azopt™	Hetlioz®	Retin-A® cream
Bethkis®	Kazano®	Symbicort®
CellCept® suspension	Kitabis® Pak	Tegretol® XR
Ciprodex®	Lialda®	Tegretol® suspension
Combigan®	Nesina®	Trileptal® suspension
Concerta®	Nexavar®	<b>Vascepa®</b>
Copaxone® 20 mg SQ	NuvaRing®	Ventolin® HFA
Daytrana®	Oseni®	<b>Viibryd®</b>
Depakote® Sprinkle	Pentasa®	Zegerid® Rx
<b>Dexilant®</b>	<b>Protonix® suspension</b>	
EpiPen	Pradaxa®	

\*\*List is subject to change

**Please keep in mind that drugs in this program may be subject to prior authorization requirements of other pharmacy programs, promoting the use of the most cost-effective product.**

## Important Billing Information

- Pursuant to this program, prescription claims submitted to the Medicaid program **do not require** the submission of Dispense as Written/Product Selection Code of '1'; **Pharmacies should submit DAW code 9** (Substitution Allowed by Prescriber but Plan Requests Brand). Pharmacies will receive an NCPDP reject response of "22" which means missing/invalid DAW code if other DAW codes are submitted. The only exception to this is DAW code 1 and "*Brand Medically Necessary*" on the prescription.
- For more information on the Brand Less Than Generic (BLTG) Program, please refer to [https://newyork.fhsc.com/providers/bltgp\\_about.asp](https://newyork.fhsc.com/providers/bltgp_about.asp)

## NYS Medicaid NYRx Mandatory Generic Drug Program

State law excludes Medicaid coverage of brand name drugs that have a Federal Food and Drug Administration (FDA) approved A-rated generic equivalent unless a prior authorization is obtained.

Coverage parameters under the Preferred Drug Program (PDP), Clinical Drug Review Program (CDRP), and/or the Brand Less Than Generic (BLTG) Program are applicable for certain products subject to the Mandatory Generic Drug Program (MGDP), including exemptions (as listed below).

### Prior Authorization Process

- Prescribers, or an agent of the prescriber, must call the prior authorization line at 1-877-309-9493 and respond to a series of questions that identify the prescriber, the patient, and the reason for prescribing this drug. The Mandatory Generic Program Prescriber Worksheet and Instructions, located at [https://newyork.fhsc.com/providers/MGDP\\_forms.asp](https://newyork.fhsc.com/providers/MGDP_forms.asp), provide step-by-step assistance in completing the prior authorization process.
- The prescriber must write “DAW and Brand Medically Necessary” on the face of the prescription.
- The call line 1-877-309-9493 is in operation 24 hours a day, seven days a week.

### Exempt Drugs

- Based on specific characteristics of the drug and/or disease state generally treated, the following brand name drugs are exempt from the program and do NOT require PA:

Exempt Drugs	
Clozaril®	Neoral®
Dilantin®	Sandimmune®
Gengraf®	Tegretol®
Lanoxin®	Zarontin®
Levothyroxine Sodium (Unithroid®, Synthroid®, Levoxyl®)	

For more information on the Mandatory Generic Program, please refer to [https://newyork.fhsc.com/providers/MGDP\\_about.asp](https://newyork.fhsc.com/providers/MGDP_about.asp).

## NYS Medicaid NYRx Dose Optimization Program

On November 14, 2013, the Medicaid NYRx program instituted a Dose Optimization initiative. Dose optimization can reduce prescription costs by reducing the number of pills a patient needs to take each day. The Department has identified drugs to be included in this program, the majority of which have FDA approval for once-a-day dosing, have multiple strengths available in correlating increments at similar costs and are currently being utilized above the recommended dosing frequency. Prior authorization will be required to obtain the following medication beyond the following limits:

### Dose Optimization Chart

Brand Name	Dose Optimization Limitations					
<b>CARDIOVASCULAR</b>						
<b>Angiotensin Receptor Blockers (ARBs)</b>						
Benicar® 20 mg	1 daily	Tablet				
Micardis® 20 mg, 40 mg	1 daily	Tablet				
Diovan® 40 mg, 80 mg, 160 mg	1 daily	Tablet				
<b>Antiarrhythmics</b>						
Amiodarone 100 mg	1 daily	Tablet	In case of dose titration for these medications, the department will allow for multiday dosing (up to 2 doses daily) for loading dose for 30 days			
<b>ARBs Combinations</b>						
Exforge® 5–160mg	1 daily	Tablet				
<b>ARBs/Diuretics</b>						
Benicar® HCT 20–12.5 mg	1 daily	Tablet				
Diovan® HCT 80–12.5 mg, 160–12.5 mg	1 daily	Tablet				
Edarbyclor® 40–12.5 mg	1 daily	Tablet				
Micardis® HCT 40–12.5 mg, 80–12.5 mg	1 daily	Tablet				
<b>Beta Blockers</b>						
Bystolic® 2.5 mg, 5 mg, 10 mg	1 daily	Tablet				
Coreg® CR 20 mg, 40 mg	1 daily	Tablet				
nadolol 40 mg	1 daily	Tablet				
Toprol® XL 25 mg, 50 mg, 100 mg	1 daily	Tablet				

Brand Name	Dose Optimization Limitations					
<b>CARDIOVASCULAR</b>						
<b>HMG Co A Reductase Inhibitors</b>						
Crestor® 5 mg, 10 mg, 20 mg	1 daily	Tablet				
<b>Niacin Derivatives</b>						
Niaspan® 500 mg	1 daily	Tablet				

Brand Name	Dose Optimization Limitations					
<b>CENTRAL NERVOUS SYSTEM</b>						
<b>Anticonvulsants</b>						
Aptiom® 200 mg, 400 mg	1 daily	Tablet				
Fycompa® 4 mg, 6 mg	1 daily	Tablet				
topiramate ER 100 mg	1 daily	Capsule				
Lamictal XR® 50 mg	1 daily	Tablet	In case of dose titration for these medications, the department will allow for multiday dosing (up to 2 doses daily) for titration purposes for 90 days			
Oxtellar XR® 300 mg	1 daily	Tablet	In case of dose titration for these medications, the department will allow for multiday dosing (up to 2 doses daily) for titration purposes for 90 days			
Lyrica® 25 mg, 50 mg, 75 mg, 100 mg, 150 mg, 200 mg	3 daily	Tablet	Electronic bypass for diagnosis of seizure disorder identified in medical claims data. In case of dose titration for these medications, the department will allow for multiday dosing (up to 2 doses daily) for titration purposes for 3 months			
Lyrica® 225 mg and 300 mg	2 daily	Tablet				
Trokendi XR® 100 mg	1 daily	Tablet				
<b>Antiparkinson Agents</b>						
Azilect® 0.5 mg	1 daily	Tablet				

Brand Name	Dose Optimization Limitations					
<b>CENTRAL NERVOUS SYSTEM</b>						
<b>Antipsychotics – Second Generation</b>						
Abilify® 2 mg	4 daily	Tablet	In case of dose titration for these medications, the Department will allow for multiday dosing (up to 2 doses/daily) for titration purposes for three months			
Abilify® 5 mg, 10 mg, 15 mg	1 daily	Tablet				
ariPIPrazole 5 mg, 10 mg, 15 mg	1 daily	Tablet				
Invega® 1.5 mg, 3 mg	1 daily	Tablet				
Latuda® 20 mg, 40 mg, 60 mg	1 daily	Tablet				
olanzapine 5 mg, 10 mg	1 daily	Tablet				
olanzapine ODT 5 mg, 10 mg	1 daily	Tablet				
paliperidone er 1.5 mg, 3 mg	1 daily	Tablet				
quetiapine fumarate er 200 mg	1 daily	Tablet				
Rexulti® 0.25 mg, 0.5 mg, 1 mg, 2 mg	1 daily	Tablet				
Seroquel® XR 150 mg, 200 mg	1 daily	Tablet				
Symbax® 3–25 mg, 6–25 mg, 12–25 mg	1 daily	Capsule				
Vraylar® 1.5 mg, 3 mg	1 daily	Capsule				
Zyprexa® Zydis 5 mg, 10 mg	1 daily	Tablet				
<b>CNS Stimulants</b>						
Adderall® XR 5 mg, 10 mg, 15 mg	1 daily	Capsule				
amphetamine salt combo ER 5 mg, 10 mg, 15 mg	1 daily	Capsule				
Concerta® ER 18 mg, 27 mg	1 daily	Tablet				
dexmethylphenidate ER 10 mg, 20 mg (Focalin XR generic)	1 daily	Capsule				
Focalin® XR 5 mg, 10 mg, 15 mg, 20 mg	1 daily	Capsule				
methylphenidate CD 10 mg, 20 mg	1 daily	Capsule				
methylphenidate er 18 mg (Concerta® generic)	1 daily	Tablet				
methylphenidate la 20 mg (Ritalin® LA generic)	1 daily	Capsule				
modafinil 100 mg	1 daily	Tablet				
Provigil® 100 mg	1 daily	Tablet				
QuilliChew® ER 20 mg	1 daily	Tablet				
Ritalin® LA 10 mg, 20 mg	1 daily	Capsule				

Brand Name	Dose Optimization Limitations		
<b>CENTRAL NERVOUS SYSTEM</b>			
Vyvanse® 10 mg, 20 mg, 30 mg, 40 mg	1 daily	Capsule	
<b>Other Agents for Attention Deficit Hyperactivity Disorder (ADHD)</b>			
guanfacine ER 1 mg, 2 mg	1 daily	Tablet	
atomoxetine 40 mg	1 daily	Capsule	
Intuniv® 1 mg, 2 mg	1 daily	Tablet	
Strattera® 40 mg	1 daily	Capsule	
<b>Sedative Hypnotics</b>			
Lunesta® 1 mg	1 daily	Tablet	
<b>Serotonin-Norepinephrine Reuptake Inhibitors (SNRIs)</b>			
Effexor® XR 37.5 mg, 75 mg, 150 mg	1 daily	Capsule	In the case of dose titration for these medications, the Department will allow for multiday dosing (up to 2 doses/daily) for titration purposes for three months.
desvenlafaxine succinate ER (Pristiq® ER 50 mg)	1 daily	Tablet	
<b>Selective Serotonin Reuptake Inhibitors (SSRIs)</b>			
Lexapro® 5 mg, 10 mg	1 daily	Tablet	In the case of dose titration for these once daily medications, the Department will allow for multiday dosing (up to 2 doses/daily) for titration purposes for three months.
Trintellix® 5 mg, 10 mg	1 daily	Tablet	
Viibryd® 10 mg, 20 mg	1 daily	Tablet	
<b>Miscellaneous Antidepressants</b>			
bupropion xl 150 mg	1 daily	Tablet	In case of dose titration for these medications, the Department will allow for multiday dosing (up to 2 doses/daily) for titration purposes for three months
mirtazapine 7.5 mg	1 daily	Tablet	

Brand Name	Dose Optimization Limitations					
<b>ENDOCRINE AND METABOLIC</b>						
<b>Biguanides</b>						
metformin ER 500 mg (Glumetza ER, Fortamet ER generic)	1 daily	Tablet				

Brand Name	Dose Optimization Limitations					
<b>ENDOCRINE AND METABOLIC</b>						
<b>Dipeptidyl Peptidase-4 (DPP-4) Inhibitors</b>						
Januvia® 25 mg, 50 mg	1 daily	Tablet				
Onglyza® 2.5 mg	1 daily	Tablet				
<b>Thiazolidinediones (TZDs)</b>						
Actos® 15 mg	1 daily	Tablet				

Brand Name	Dose Optimization Limitations					
<b>GASTROINTESTINAL</b>						
<b>Proton Pump Inhibitors</b>						
Dexilant® 30 mg	1 daily	Capsule				
Nexium® 5 mg, 10 mg, 20 mg	1 daily	Packet				
Nexium® 20 mg	1 daily	Capsule				
Prevacid® DR 15 mg	1 daily	Capsule				

Brand Name	Dose Optimization Limitations					
<b>HEMATOLOGICAL</b>						
<b>Anticoagulants - Oral</b>						
Xarelto® 10 mg	1 daily	Capsule				

Brand Name	Dose Optimization Limitations					
<b>RENAL AND GENITOURINARY</b>						
<b>Urinary Tract Antispasmodics</b>						
Detrol® LA 2 mg	1 daily	Capsule				
Myrbetriq® 25 mg	1 daily	Tablet				
oxybutynin chloride ER 5 mg	1 daily	Tablet				

Brand Name	Dose Optimization Limitations		
<b>RENAL AND GENITOURINARY</b>			
Toviaz® ER 4 mg	1 daily	Tablet	
VESIcare® 5 mg	1 daily	Tablet	

PA requirements are not dependent on the date a prescription is written. New prescriptions and refills on existing prescriptions require PA even if the prescription was written before the date the drug was determined to require PA.

To obtain a prior authorization (PA), please call the prior authorization Clinical Call Center at 1-877-309-9493. The Clinical Call Center is available 24 hours per day, 7 days per week with pharmacy technicians and pharmacists who will work with you, or your agent, to quickly obtain PA.

Medicaid enrolled prescribers with an active e-PACES account can initiate PA requests through the web-based application PAXpress®. The website for PAXpress is <https://paxpress.nypa.hidinc.com>.

When, in the judgment of the prescriber or the pharmacist, an emergency condition exists, the prescriber or pharmacist can call the Clinical Call center and obtain authorization for a 72-hour emergency supply of the drug prescribed to allow time for the prior authorization to be obtained.